

Cauchy-Riemann Equations and Conformal Mapping

26.2



Introduction

In this Section we consider two important features of complex functions. The Cauchy Riemann equations introduced on page 2 provide a necessary and sufficient condition for a function $f(z)$ to be analytic in some region of the complex plane; this allows us to find $f'(z)$ in that region by the rules of the previous Section. A mapping between the z -plane and the w -plane is said to be conformal if the angle between two intersecting curves in the z -plane is equal to the angle between their mappings in the w -plane. Such a mapping has widespread uses in solving problems in fluid flow and electromagnetics, for example, where the given problem geometry is somewhat complicated.



Prerequisites

Before starting this Section you should ...

- ① understand the idea of a complex function and its derivative



Learning Outcomes

After completing this Section you should be able to ...

- ✓ use the Cauchy Riemann equations to obtain the derivative of complex functions
- ✓ appreciate the idea of a conformal mapping

1. Cauchy-Riemann equations

Remembering that $z = x + iy$ and $w = u + iv$ we note that there is a very useful test to determine whether a function $w = f(z)$ is analytic at a point. This is provided by the **Cauchy-Riemann** equations. These state that $w = f(z)$ is differentiable at a point $z = z_0$ if, and only if,

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \text{and} \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x} \quad \text{at that point.}$$

When these equations hold then it can be shown that the complex derivative may be determined by using either $\frac{df}{dz} = \frac{\partial f}{\partial x}$ or $\frac{df}{dz} = -i\frac{\partial f}{\partial y}$.

(The use of ‘if, and only if,’ means that if the equations are valid, then the function is differentiable **and vice versa**).

If we consider $f(z) = z^2 = x^2 - y^2 + 2ixy$ then $u = x^2 - y^2$ and $v = 2xy$ so that

$$\frac{\partial u}{\partial x} = 2x, \quad \frac{\partial u}{\partial y} = -2y, \quad \frac{\partial v}{\partial x} = 2y, \quad \frac{\partial v}{\partial y} = 2x.$$

It should be clear that, for this example, the Cauchy-Riemann equations are always satisfied; therefore, the function is analytic everywhere. We find that

$$\frac{df}{dz} = \frac{\partial f}{\partial x} = 2x + 2iy = 2z \quad \text{or, equivalently} \quad \frac{df}{dz} = -i\frac{\partial f}{\partial y} = -i(-2y + 2ix) = 2z$$

This is the result we would expect to get by simply differentiating $f(z)$ as if it was a real function. For analytic functions **this will always be the case**.

Example Show that the function $f(z) = z^3$ is analytic everywhere and hence obtain its derivative.

Solution

$$w = f(z) = (x + iy)^3 = x^3 - 3xy^2 + (3x^2y - y^3)i$$

Hence

$$u = x^3 - 3xy^2 \quad \text{and} \quad v = 3x^2y - y^3.$$

Then

$$\frac{\partial u}{\partial x} = 3x^2 - 3y^2, \quad \frac{\partial u}{\partial y} = -6xy, \quad \frac{\partial v}{\partial x} = 6xy, \quad \frac{\partial v}{\partial y} = 3x^2 - 3y^2.$$

The Cauchy-Riemann equations are identically true and $f(z)$ is analytic everywhere.

Furthermore

$$\frac{df}{dz} = \frac{\partial f}{\partial x} = 3x^2 - 3y^2 + (6xy)i = 3(x + iy)^2 = 3z^2 \quad \text{as we would expect.}$$

We can easily find functions which are not analytic anywhere and others which are only analytic in a restricted region of the complex plane. Consider again the function $f(z) = \bar{z} = x - iy$. Here

$$u = x \quad \text{so that} \quad \frac{\partial u}{\partial x} = 1, \quad \text{and} \quad \frac{\partial u}{\partial y} = 0; \quad v = -y \quad \text{so that} \quad \frac{\partial v}{\partial x} = 0, \quad \frac{\partial v}{\partial y} = -1.$$

The Cauchy-Riemann equations are never satisfied so that \bar{z} is not differentiable anywhere and so is not analytic anywhere.

By contrast if we consider the function $f(z) = \frac{1}{z}$ we find that

$$u = \frac{x}{x^2 + y^2}; \quad v = \frac{y}{x^2 + y^2}.$$

As can readily be shown, the Cauchy-Riemann equations are satisfied everywhere except for $x^2 + y^2 = 0$, i.e. $x = y = 0$ (or, equivalently, $z = 0$). At all other points $f'(z) = -\frac{1}{z^2}$. This function is analytic everywhere except at the single point $z = 0$.

Analyticity is a very powerful property of a function of a complex variable. Such functions tend to behave like functions of a real variable.

Example Show that if $f(z) = z\bar{z}$ then $f'(z)$ exists only at $z = 0$.

Solution

$$f(z) = x^2 + y^2 \quad \text{so that} \quad u = x^2 + y^2, \quad v = 0.$$

$$\frac{\partial u}{\partial x} = 2x, \quad \frac{\partial u}{\partial y} = 2y, \quad \frac{\partial v}{\partial x} = 0, \quad \frac{\partial v}{\partial y} = 0.$$

Hence the Cauchy-Riemann equations are satisfied only where $x = 0$ and $y = 0$, i.e. where $z = 0$. Therefore this function is not analytic anywhere.

Analytic Functions and Harmonic Functions

Using the Cauchy-Riemann equations in a region of the z -plane where $f(z)$ is analytic,

$$\frac{\partial^2 u}{\partial x \partial y} = \frac{\partial}{\partial x} \left(\frac{\partial u}{\partial y} \right) = \frac{\partial}{\partial x} \left(-\frac{\partial v}{\partial x} \right) = -\frac{\partial^2 v}{\partial x^2}$$

and

$$\frac{\partial^2 u}{\partial y \partial x} = \frac{\partial}{\partial y} \left(\frac{\partial u}{\partial x} \right) = \frac{\partial}{\partial y} \left(\frac{\partial v}{\partial y} \right) = \frac{\partial^2 v}{\partial y^2}.$$

If these differentiations are possible then $\frac{\partial^2 u}{\partial x \partial y} = \frac{\partial^2 u}{\partial y \partial x}$ so that

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \quad (\text{Laplace's equation}).$$

In a similar way we find that

$$\frac{\partial^2 v}{\partial x^2} + \frac{\partial^2 v}{\partial y^2} = 0 \quad (\text{can you show this?})$$

When $f(z)$ is analytic the functions u and v are called **conjugate harmonic functions**.

Suppose $u = u(x, y) = xy$ then it is easy to verify that u satisfies Laplace's equation (try this). We now try to find the conjugate harmonic function $v = v(x, y)$.

First, using the Cauchy-Riemann equations:

$$\frac{\partial v}{\partial y} = \frac{\partial u}{\partial x} = y \quad \text{and} \quad \frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y} = -x.$$

Integrating the first equation gives $v = \frac{1}{2}y^2 +$ a function of x . Integrating the second equation gives $v = -\frac{1}{2}x^2 +$ a function of y . Bearing in mind that an additive constant leaves no trace after differentiation we pool the information above to obtain

$$v = \frac{1}{2}(y^2 - x^2) + C \quad \text{where } C \text{ is a constant}$$

Note that $f(z) = u + iv = xy + \frac{1}{2}(y^2 - x^2)i + D$ where D is a constant (replacing Ci).

We can write $f(z) = -\frac{1}{2}iz^2 + D$ (as you can verify). This function is analytic everywhere.



Show that the function $u = x^2 - x - y^2$ is harmonic.

Your solution

Hence $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$ and u is harmonic.

$$\frac{\partial u}{\partial x} = 2x - 1, \quad \frac{\partial^2 u}{\partial x^2} = 2, \quad \frac{\partial u}{\partial y} = -2y, \quad \frac{\partial^2 u}{\partial y^2} = -2.$$



Now find the conjugate harmonic function v .

Your solution

Integrating $\frac{\partial v}{\partial x} = 2x - y + C$ gives $v = x^2 - xy + Cx + D$, where C is a constant. Ignoring the duplication, $v = 2xy - y + C$, where C is a constant.



Finally, find $f(z)$ in terms of z .

Your solution

Now $z^2 - y^2 + 2ixy$ and $z = x + iy$ thus $f(z) = z^2 - z + D$.
 $f(z) = u + iv = x^2 - y^2 + 2ixy - iy + D$, D constant

Exercises

- Find the singular point of the rational function $f(z) = \frac{z}{z - 2i}$. Find $f'(z)$ at other points and evaluate $f'(-i)$.
- Show that the function $f(z) = z^2 + z$ is analytic everywhere and hence obtain its derivative.
- Show that the function $u = x^2 - y^2 - 2y$ is harmonic, find the conjugate harmonic function v and hence find $f(z) = u + iv$ in terms of z .

$$\frac{df}{dz} = \frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} = 2x + 1 + 2iy + i$$

Here the Cauchy-Riemann equations are identically true and $f(z)$ is analytic everywhere.

$$\frac{\partial u}{\partial x} = 2x + 1, \quad \frac{\partial u}{\partial y} = -2y, \quad \frac{\partial v}{\partial x} = 2y, \quad \frac{\partial v}{\partial y} = 2x + 1$$

$$2. \quad u = x^2 - y^2 + x \quad \text{and} \quad v = 2xy + y$$

$$f'(-i) = \frac{-2!}{-2!} \frac{(-3i)^2}{-9} = \frac{9}{2}$$

$$f'(z) = \frac{(z - 2i)^{-2}}{(z - 2i)^{-2}} = \frac{1}{z - 2i}$$

1. $f(z)$ is singular at $z = 2i$. Elsewhere

$$\begin{aligned}
 & \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \quad \text{therefore } u \text{ is harmonic.} \\
 & \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \text{therefore } v = \int \frac{\partial u}{\partial x} dy + f(x) \\
 & \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x} \quad \text{therefore } v = -\int \frac{\partial u}{\partial y} dx + g(y)
 \end{aligned}$$

2. Conformal Mapping

In Section 1 we saw that the real and imaginary parts of an analytic function each satisfies Laplace's equation. We shall show now that the curves

$$u(x, y) = \text{constant} \quad \text{and} \quad v(x, y) = \text{constant}$$

intersect each other at right angles (we say that they are *orthogonal*). To see this we note that along the curve $u(x, y) = \text{constant}$ we have $du = 0$. Hence

$$du = \frac{\partial u}{\partial x} dx + \frac{\partial u}{\partial y} dy = 0.$$

Thus, on these curves the gradient at a general point is given by

$$\frac{dy}{dx} = -\frac{\frac{\partial u}{\partial x}}{\frac{\partial u}{\partial y}}.$$

Similarly along the curve $v(x, y) = \text{constant}$, we have

$$\frac{dy}{dx} = -\frac{\frac{\partial v}{\partial x}}{\frac{\partial v}{\partial y}}.$$

The product of these gradients is

$$\frac{(\frac{\partial u}{\partial x})(\frac{\partial v}{\partial x})}{(\frac{\partial u}{\partial y})(\frac{\partial v}{\partial y})} = -\frac{(\frac{\partial u}{\partial x})(\frac{\partial u}{\partial y})}{(\frac{\partial u}{\partial y})(\frac{\partial u}{\partial x})} = -1$$

where we have made use of the Cauchy-Riemann equations. We deduce that the curves are orthogonal.

As an example of the practical application of this work consider two-dimensional electrostatics. If $u = \text{constant}$ gives the *equipotential* curves then the curves $v = \text{constant}$ are the *electric lines of force*. Figure 1 shows some curves from each set in the case of oppositely-charged particles near

to each other; the dashed curves are the lines of force and the solid curves are the equipotentials.




Figure 1

In ideal fluid flow the curves $v = \text{constant}$ are the *streamlines* of the flow.

In these situations the function $w = u + iv$ is the **complex potential** of the field.

Conformal Mapping

A function $w = f(z)$ can be regarded as a mapping, which ‘maps’ a point in the z -plane to a point in the w -plane. Curves in the z -plane will be mapped into curves in the w -plane.

Consider aerodynamics. The idea is that we are interested in the fluid flow, in a complicated geometry (say flow past an aerofoil). We first find the flow in a simple geometry that can be mapped to the aerofoil shape (the complex plane with a circular hole works here). Most of the calculations necessary to find physical characteristics such as lift and drag on the aerofoil can be performed in the simple geometry - the resulting integrals being much easier to evaluate than in the complicated geometry.

Consider the mapping

$$w = z^2.$$

The point $z = 2 + i$ maps to $w = (2 + i)^2 = 3 + 4i$. The point $z = 2 + i$ lies on the intersection of the two lines $x = 2$ and $y = 1$. To what curves do these map? To answer this question we note that a point on the line $y = 1$ can be written as $z = x + i$. Then

$$w = (x + i)^2 = x^2 - 1 + 2xi$$

As usual, let $w = u + iv$, then

$$u = x^2 - 1 \quad \text{and} \quad v = 2x$$

Eliminating x we obtain:

$$4u = 4x^2 - 4 = v^2 = 4 \quad \text{or} \quad v^2 = 4 + 4u.$$

Example To what curve does the line $x = 2$ map?

Solution

A point on the line is $z = 2 + yi$. Then

$$w = (2 + yi)^2 = 4 - y^2 + 4yi$$

Hence $u = 4 - y^2$ and $v = 4y$ so that, eliminating y we obtain

$$16u = 64 - v^2 \quad \text{or} \quad v^2 = 64 - 16u$$

In Figure 2(a) we sketch the lines $x = 2$ and $y = 1$ and in Figure 2(b) we sketch the curves into which they map. Note these curves intersect at the point $(3,4)$.




Figure 2

The angle between the original lines was clearly 90° ; what is the angle between the curves at the point of intersection?

The curve $v^2 = 4 + 4u$ has a gradient $\frac{dv}{du}$. Differentiating the equation implicitly we obtain

$$2v \frac{dv}{du} = 4 \quad \text{or} \quad \frac{dv}{du} = \frac{2}{v}$$

At the point $(3,4)$ $\frac{dv}{du} = \frac{1}{2}$.



Find $\frac{dv}{du}$ for the curve $v^2 = 64 - 16u$ and evaluate it at the point $(3,4)$

Your solution

$$\frac{dv}{dz} = -\frac{u}{v} \quad \text{At } v = 4 \text{ we obtain } \frac{du}{dz} = -2$$

$$\frac{dv}{dz} = -\frac{u}{v} \quad \therefore \quad \frac{du}{dz} = -\frac{u}{v}$$

Note that the product of the gradients at $(3,4)$ is -1 and therefore the angle between the curves at their point of intersection is also 90° . Since the angle between the lines and the angle between the curves is the same we say the angle is preserved.

In general, if two curves in the z -plane intersect at a point z_0 , and their image curves under the mapping $w = f(z)$ intersect at $w_0 = f(z_0)$ and the angle between the two original curves at z_0 equals the angle between the image curves at w_0 we say that the mapping is *conformal* at z_0 .

An analytic function is conformal everywhere except where $f'(z) = 0$.



At which points is $w = e^z$ not conformal?

Your solution

$f'(z) = e^z$. Since this is never zero the mapping is conformal everywhere.

Inversion

The mapping

$$w = f(z) = \frac{1}{z}$$

is called an *inversion*. It maps the interior of the unit circle in the z -plane to the exterior of the unit circle in the w -plane, and vice-versa. Note that

$$w = u + iv = \frac{x}{x^2 + y^2} - \frac{y}{x^2 + y^2}i \quad \text{and similarly} \quad z = x + iy = \frac{u}{u^2 + v^2} - \frac{v}{u^2 + v^2}i$$

so that

$$u = \frac{x}{x^2 + y^2} \quad \text{and} \quad v = -\frac{y}{x^2 + y^2}.$$

A line through the origin in the z -plane will be mapped into a line through the origin in the w -plane. To see this consider the line $y = mx$, for m constant. Then

$$u = \frac{x}{x^2 + m^2x^2} \quad \text{and} \quad v = -\frac{mx}{x^2 + m^2x^2}$$

so that $v = -mu$, which is a line through the origin in the w -plane.



Consider the line $ax + by + c = 0$ where $c \neq 0$. This represents a line in the z -plane which does not pass through the origin. To what sort of curve does it map in the w -plane?

Your solution

which is the equation of a circle in the w -plane which passes through the origin.

$$0 = \frac{c}{a} + \frac{c}{b} + v^2 + u^2$$

Hence $au - bv + c(u^2 + v^2) = 0$. Dividing by c we obtain the equation:

$$0 = \frac{u^2 + v^2}{a} - \frac{u^2 + v^2}{b} + c = 0$$

The mapped curve is

Similarly, it can be shown that a circle in the z -plane passing through the origin maps to a line in the w -plane which does not pass through the origin and a circle in the z -plane which does not pass through the origin maps to a circle in the w -plane which does not pass through the origin. The inversion mapping is an example of the *bilinear transformation*:

$$w = f(z) = \frac{az + b}{cz + d} \quad \text{where we demand that } ad - bc \neq 0$$

(If $ad - bc = 0$ the mapping reduces to $f(z) = \text{constant}$).



Find the bilinear transformations which map $z = 2$ to $w = 1$.

Your solution

$$1 = \frac{2a + b}{2c + d} \text{ Hence } 2a + b = 2c + d$$



If in addition, $z = -1$ is mapped to $w = 3$ find the class of transformation which is possible.

Your solution

$$3 = \frac{p + c - a + v - q}{q + v - a + b} \text{ Hence } p + c - a + v - q = 3(q + v - a + b)$$

If, further, $z = 0$ is mapped to $w = -5$ then $-5 = \frac{b}{d}$ so that $b = -5d$. Substituting this last relation into the first two obtained we obtain

$$\begin{aligned} 2a - 2c - 6d &= 0 \\ -a + 3c - 8d &= 0 \end{aligned}$$

Solving these two in terms of d we find $2c = 11d$ and $2a = 17d$. Hence the transformation is:

$$w = \frac{17z - 10}{11z + 2} \text{ (note that the } d\text{'s cancel in the numerator and denominator).}$$

Some other mappings are shown in Figure 3.




Figure 3

As an engineering application we consider the transformation

$$w = z - \frac{\ell^2}{z} \quad \text{where } \ell \text{ is a constant.}$$

It is used to map circles which contain $z = 1$ as an interior point and which pass through $z = -1$ into shapes resembling aerofoils. Figure 4 shows an example




Figure 4

This creates a cusp at which the associated fluid velocity can be infinite. This can be avoided by adjusting the fluid flow in the z -plane. Eventually, this can be used to find the lift generated by such an aerofoil in terms of physical characteristics such as aerofoil shape and air density and speed.

Exercises

1. Find a bilinear transformation which maps $z = 0, -1, -i$ into $w = i, 0, 1$ respectively.

$$w = \frac{az + b}{cz + d} \quad 1.$$

$z = 0, w = i$ gives $i = \frac{b}{d}$ so that $b = di$

$z = -1, w = 1$ gives $1 = \frac{-a + b}{-c + d}$ so that $-ci + d = -a + di + d$

$z = -i, w = 1$ gives $1 = \frac{-ai + b}{-ci + d}$ so that $-ci + d = -ai + d + di$

Hence $c = -d$ and $m = \frac{d + zp}{di + d} = \frac{1 + z -}{1 + i}$