

Lines and Planes

9.5



Introduction

Direction ratios provide a convenient way of specifying the direction of a line in three dimensional space. Direction cosines are the cosines of the angles between a line and the coordinate axes. In this Section we show how these quantities are calculated.

Vectors are very convenient tools for analysing lines and planes in three dimensions. In this Section you will learn how to formulate the vector equation of a line and the vector equation of a plane.



Prerequisites

Before starting this Section you should ...

- ① understand and be able to calculate the scalar product of two vectors
- ② understand and be able to calculate the vector product of two vectors



Learning Outcomes

After completing this Section you should be able to ...

- ✓ obtain the vector equation of a line
- ✓ obtain the vector equation of a plane passing through a given point and which is perpendicular to a given vector
- ✓ obtain the vector equation of a plane which is a given distance from the origin and which is perpendicular to a given vector

1. The Direction Ratio and Direction Cosines

Consider the point $P(4, 5)$ and its position vector $4\underline{i} + 5\underline{j}$ shown in Figure 1.




Figure 1.

The **direction ratio** of the vector \overrightarrow{OP} is defined to be 4:5. We can interpret this as stating that to move in the direction of the line OP we must move 4 units in the x direction for every 5 units in the y direction.

The **direction cosines** of the vector \overrightarrow{OP} are the cosines of the angles between the vector and each of the axes. Specifically, referring to Figure 1 these are

$$\cos \alpha \quad \text{and} \quad \cos \beta$$

Noting that the length of \overrightarrow{OP} is $\sqrt{4^2 + 5^2} = \sqrt{41}$ we can write

$$\cos \alpha = \frac{4}{\sqrt{41}}, \quad \cos \beta = \frac{5}{\sqrt{41}}$$

It is conventional to label the direction cosines as ℓ and m so that

$$\ell = \frac{4}{\sqrt{41}}, \quad m = \frac{5}{\sqrt{41}}$$

More generally we have the following result:



Key Point

For *any* vector $\underline{r} = a\underline{i} + b\underline{j}$, its direction ratio is $a : b$. Its direction cosines are

$$\ell = \frac{a}{\sqrt{a^2 + b^2}}, \quad m = \frac{b}{\sqrt{a^2 + b^2}}$$

Example Point A has coordinates $(3, 5)$, and point B has coordinates $(7, 8)$.

- a) Write down the vector \overrightarrow{AB} . b) Find the direction ratio of the vector \overrightarrow{AB} .
 c) Find its direction cosines, ℓ and m . d) Show that $\ell^2 + m^2 = 1$

Solution

a) $\overrightarrow{AB} = \underline{b} - \underline{a} = 4\underline{i} + 3\underline{j}$. b) The direction ratio of \overrightarrow{AB} is therefore 4:3. c) The direction cosines are

$$\ell = \frac{4}{\sqrt{4^2 + 3^2}} = \frac{4}{5}, \quad m = \frac{3}{\sqrt{4^2 + 3^2}} = \frac{3}{5}$$

d)

$$\ell^2 + m^2 = \left(\frac{4}{5}\right)^2 + \left(\frac{3}{5}\right)^2 = \frac{16}{25} + \frac{9}{25} = \frac{25}{25} = 1$$

The final result in the previous example is true in general:



Key Point

If ℓ and m are the direction cosines of a vector lying in the xy plane, then $\ell^2 + m^2 = 1$

Exercises

1. P and Q have coordinates $(-2, 4)$ and $(7, 8)$ respectively.
 a) Find the direction ratio of the vector \overrightarrow{PQ} b) Find the direction cosines of \overrightarrow{PQ} .

Answers 1. a) 9 : 4, b) $\frac{\sqrt{97}}{4}, \frac{\sqrt{97}}{9}$.

2. Direction Ratios and Cosines in Three Dimensions

The concepts of direction ratio and direction cosines extend naturally to three dimensions. Consider Figure 2.

Given a vector $\underline{r} = a\underline{i} + b\underline{j} + c\underline{k}$ its direction ratios are $a : b : c$. This means that to move in the direction of the vector we must move a units in the x direction and b units in the y direction for every c units in the z direction.

The direction cosines are the cosines of the angles between the vector and each of the axes. It is conventional to label direction cosines as ℓ , m and n and they are given by

$$\ell = \cos \alpha = \frac{a}{\sqrt{a^2 + b^2 + c^2}}, \quad m = \cos \beta = \frac{b}{\sqrt{a^2 + b^2 + c^2}}, \quad n = \cos \gamma = \frac{c}{\sqrt{a^2 + b^2 + c^2}}$$

In general we have the following result:




Figure 2.



Key Point

For any vector $\underline{r} = a\underline{i} + b\underline{j} + c\underline{k}$ its direction ratios are $a : b : c$. Its direction cosines are

$$\ell = \frac{a}{\sqrt{a^2 + b^2 + c^2}}, \quad m = \frac{b}{\sqrt{a^2 + b^2 + c^2}}, \quad n = \frac{c}{\sqrt{a^2 + b^2 + c^2}}$$

where $\ell^2 + m^2 + n^2 = 1$

Exercises

- Points A and B have position vectors $\underline{a} = -3\underline{i} + 2\underline{j} + 7\underline{k}$, and $\underline{b} = 3\underline{i} + 4\underline{j} - 5\underline{k}$ respectively. Find
 - \overrightarrow{AB}
 - $|\overrightarrow{AB}|$
 - the direction ratios of \overrightarrow{AB}
 - the direction cosines (ℓ, m, n) of \overrightarrow{AB} .
 - Show that $\ell^2 + m^2 + n^2 = 1$.
- Find the direction ratios, the direction cosines and the angles that the vector \overrightarrow{OP} makes with each of the axes when P is the point with coordinates $(2, 4, 3)$.
- A line is inclined at 60° to the x axis and 45° to the y axis. Find its inclination to the z axis.

Answers 1. a) $6\underline{i} + 2\underline{j} - 12\underline{k}$, b) $\sqrt{184}$, c) $6 : 2 : -12$, d) $\frac{\sqrt{184}}{6}, \frac{\sqrt{184}}{2}, \frac{\sqrt{184}}{-12}$, e) $2 : 2 : 4 : 3; \frac{\sqrt{29}}{2}, \frac{\sqrt{29}}{3}, \frac{\sqrt{29}}{4}; 68.2^\circ, 42.0^\circ, 56.1^\circ, 3.60^\circ$ or 120° .

3. The Vector Equation of a Line

Consider the straight line APB shown in Figure 3. This is a line in three-dimensional space.




Figure 3.

Points A and B are fixed and known points on the line, and have position vectors \underline{a} and \underline{b} respectively. Point P is any other arbitrary point on the line, and has position vector \underline{r} . Note that because \overrightarrow{AB} and \overrightarrow{AP} are parallel, \overrightarrow{AP} is simply a scalar multiple of \overrightarrow{AB} , that is, $\overrightarrow{AP} = t\overrightarrow{AB}$ where t is a number.



Write down an expression for the vector \overrightarrow{AB} in terms of \underline{a} and \underline{b} .

Your solution

$$\underline{b} - \underline{a} = \overrightarrow{AB}$$



Use the triangle law for vector addition to find an expression for \underline{r} in terms of \underline{a} , \underline{b} and t .

Your solution

$$\overrightarrow{AP} = t\overrightarrow{AB} \quad \text{since} \quad (\overline{p} - \overline{q})t + \overline{p} = \overline{x}$$

so that

$$\overrightarrow{OP} + \overrightarrow{AO} = \overrightarrow{AO}$$

The answer to the above exercise $\underline{r} = \underline{a} + t(\underline{b} - \underline{a})$ is the **vector equation of the line** through A and B . It is a rule which gives the position vector \underline{r} of a general point on the line in terms of the **given** vectors $\underline{a}, \underline{b}$. By varying the value of t we can move to any point on the line. For example, when

$$t = 0, \quad \text{the equation gives} \quad \underline{r} = \underline{a}, \quad \text{which locates point } A$$

$$t = 1, \quad \text{the equation gives} \quad \underline{r} = \underline{b}, \quad \text{which locates point } B$$

If $0 < t < 1$ the point P lies on the line between A and B . If $t > 1$ the point P lies on the line beyond B . If $t < 0$ the point P lies on the line beyond A to the left of the figure.



Key Point

The **vector equation of the line** through points A and B with position vectors \underline{a} and \underline{b} is

$$\underline{r} = \underline{a} + t(\underline{b} - \underline{a})$$



Write down the vector equation of the line which passes through the points with position vectors $\underline{a} = 3\underline{i} + 2\underline{j}$ and $\underline{b} = 7\underline{i} + 5\underline{j}$.

Your solution

$$\begin{pmatrix} 3 \\ 2 \\ 4 \end{pmatrix} t + \begin{pmatrix} 2 \\ 3 \\ 1 \end{pmatrix} = \underline{x}$$

Using column vector notation we could write

$$\begin{pmatrix} 3 \\ 2 \\ 4 \end{pmatrix} t + \begin{pmatrix} 2 \\ 3 \\ 1 \end{pmatrix} = \underline{v} - \underline{q}$$

The equation of the line is then

$$\begin{pmatrix} 3 \\ 2 \\ 4 \end{pmatrix} t + \begin{pmatrix} 2 \\ 3 \\ 1 \end{pmatrix} = \underline{v} - \underline{q}$$



Write down the vector equation of the line which passes through the points with position vectors $\underline{a} = 5\underline{i} - 2\underline{j} + 3\underline{k}$ and $\underline{b} = 2\underline{i} + \underline{j} - 4\underline{k}$.

Your solution

$$\begin{pmatrix} 2 \\ 3 \\ -3 \end{pmatrix} t + \begin{pmatrix} 3 \\ 2 \\ 5 \end{pmatrix} = \underline{v} - \underline{q}$$

The equation of the line is then

$$\begin{pmatrix} 2 \\ 3 \\ -3 \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ 5 \end{pmatrix} - \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix} = \underline{v} - \underline{q}$$

Using column vector notation note that

Cartesian form

On occasions it is useful to convert the vector form of the equation of a straight line into cartesian form. Suppose we write

$$\underline{a} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}, \quad \underline{b} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}, \quad \underline{r} = \begin{pmatrix} x \\ y \\ z \end{pmatrix}$$

then $\underline{r} = \underline{a} + t(\underline{b} - \underline{a})$ implies

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} + t \begin{pmatrix} b_1 - a_1 \\ b_2 - a_2 \\ b_3 - a_3 \end{pmatrix} = \begin{pmatrix} a_1 + t(b_1 - a_1) \\ a_2 + t(b_2 - a_2) \\ a_3 + t(b_3 - a_3) \end{pmatrix}$$

Equating the individual components we find

$$x = a_1 + t(b_1 - a_1), \quad \text{or equivalently } t = \frac{x - a_1}{b_1 - a_1}$$

$$y = a_2 + t(b_2 - a_2), \quad \text{or equivalently } t = \frac{y - a_2}{b_2 - a_2}$$

$$z = a_3 + t(b_3 - a_3), \quad \text{or equivalently } t = \frac{z - a_3}{b_3 - a_3}$$

Each expression on the right is equal to t and so we can write

$$\frac{x - a_1}{b_1 - a_1} = \frac{y - a_2}{b_2 - a_2} = \frac{z - a_3}{b_3 - a_3}$$

This gives the **cartesian form** of the equations of the straight line which passes through the points with coordinates (a_1, a_2, a_3) and (b_1, b_2, b_3) .



Key Point

The **cartesian form** of the equation of the straight line which passes through the points with coordinates (a_1, a_2, a_3) and (b_1, b_2, b_3) is

$$\frac{x - a_1}{b_1 - a_1} = \frac{y - a_2}{b_2 - a_2} = \frac{z - a_3}{b_3 - a_3}$$

- Example** a) Write down the cartesian form of the equation of the straight line which passes through the two points $(9, 3, -2)$ and $(4, 5, -1)$.
b) State the equivalent vector equation.

Solution

a)

$$\frac{x-9}{4-9} = \frac{y-3}{5-3} = \frac{z-(-2)}{-1-(-2)}$$

that is

$$\frac{x-9}{-5} = \frac{y-3}{2} = \frac{z+2}{1} \quad (\text{cartesian form})$$

The vector equation is

$$\begin{aligned} \underline{r} &= \underline{a} + t(\underline{b} - \underline{a}) \\ &= \begin{pmatrix} 9 \\ 3 \\ -2 \end{pmatrix} + t \left(\begin{pmatrix} 4 \\ 5 \\ -1 \end{pmatrix} - \begin{pmatrix} 9 \\ 3 \\ -2 \end{pmatrix} \right) \\ &= \begin{pmatrix} 9 \\ 3 \\ -2 \end{pmatrix} + t \begin{pmatrix} -5 \\ 2 \\ 1 \end{pmatrix} \end{aligned}$$

Exercises

- 1.a) Write down the vector \overrightarrow{AB} joining the points A and B with coordinates $(3, 2, 7)$ and $(-1, 2, 3)$ respectively.
b) Find the equation of the straight line through A and B .
2. Write down the vector equation of the line passing through the points with position vectors $\underline{p} = 3\underline{i} + 7\underline{j} - 2\underline{k}$ and $\underline{q} = -3\underline{i} + 2\underline{j} + 2\underline{k}$. Find also the cartesian equation of this line.
3. Find the vector equation of the line passing through $(9, 1, 2)$ and which is parallel to the vector $(1, 1, 1)$.

Answers 1. a) $-4\underline{i} - 4\underline{j} - 4\underline{k}$. b) $\underline{r} = \begin{pmatrix} 3 \\ 2 \\ 7 \end{pmatrix} + t \begin{pmatrix} -4 \\ 0 \\ -4 \end{pmatrix}$. 2. $\underline{r} = \begin{pmatrix} 3 \\ 7 \\ -2 \end{pmatrix} + t \begin{pmatrix} -6 \\ -5 \\ 4 \end{pmatrix}$. 3. $\underline{r} = \begin{pmatrix} 9 \\ 1 \\ 1 \end{pmatrix} + t \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$. Cartesian form $\frac{x-3}{-6} = \frac{y-2}{-5} = \frac{z-7}{4}$.

4. The Vector Equation of a Plane

Consider the plane shown in Figure 4. Suppose that A is a known point in the plane and has position vector \underline{a} . Suppose that P is any other arbitrary point in the plane with position vector \underline{r} . Clearly the vector \overrightarrow{AP} lies in the plane.




Figure 4.



Referring to Figure 4, find the vector \overrightarrow{AP} in terms of \underline{a} and \underline{r} .

Your solution

$$\underline{d} - \underline{r}$$

Also shown in Figure 4 is a vector which is perpendicular to the plane and denoted by \underline{n} .



What relationship exists between \underline{n} and the vector \overrightarrow{AP} ?

Your solution

Hint: think about the scalar product.

$$\underline{r} \cdot \underline{n} = \underline{a} \cdot \underline{n}$$

or alternatively $\underline{r} \cdot \underline{n} - \underline{a} \cdot \underline{n} = 0$ so that

$$0 = \underline{n} \cdot (\underline{r} - \underline{a})$$

Because \overrightarrow{AP} and \underline{n} are perpendicular their scalar product must equal zero, that is

The answer to the above exercise $\underline{r} \cdot \underline{n} = \underline{a} \cdot \underline{n}$ is the **equation of a plane**, written in vector form, passing through A and perpendicular to \underline{n} .



Key Point

A plane passing through the point with position vector \underline{a} , and perpendicular to the vector \underline{n} has equation

$$\underline{r} \cdot \underline{n} = \underline{a} \cdot \underline{n}$$

In this formula it does not matter whether or not \underline{n} is a unit vector. If $\hat{\underline{n}}$ is a unit vector then $\underline{a} \cdot \hat{\underline{n}}$ represents the perpendicular distance from the origin to the plane which we usually denote by d (for details of this see Section 9.3). Hence we can write

$$\underline{r} \cdot \hat{\underline{n}} = d$$

This is the **equation of a plane**, written in vector form, with unit normal $\hat{\underline{n}}$ and which is a perpendicular distance d from O .



Key Point

A plane with unit normal $\hat{\underline{n}}$, which is a perpendicular distance d from O is given by

$$\underline{r} \cdot \hat{\underline{n}} = d$$

- Example**
- Find the vector equation of the plane which passes through the point with position vector $3\underline{i} + 2\underline{j} + 5\underline{k}$ and which is perpendicular to $\underline{i} + \underline{k}$.
 - Find the cartesian equation of this plane.

Solution

a) Using the previous results we can write down the equation

$$\underline{r} \cdot (\underline{i} + \underline{k}) = (3\underline{i} + 2\underline{j} + 5\underline{k}) \cdot (\underline{i} + \underline{k}) = 3 + 5 = 8$$

b) Writing \underline{r} as $x\underline{i} + y\underline{j} + z\underline{k}$ we have the cartesian form:

$$(x\underline{i} + y\underline{j} + z\underline{k}) \cdot (\underline{i} + \underline{k}) = 8$$

so that

$$x + z = 8$$



- a) Find the vector equation of the plane through $(7, 3, -5)$ for which $\underline{n} = (1, 1, 1)$ is a vector normal to the plane.
 b) What is the distance of the plane from O?

Your solution

$$\frac{8}{5} = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \cdot \frac{1}{5} \begin{pmatrix} 7 \\ 3 \\ -5 \end{pmatrix} = \underline{n} \cdot \underline{p}$$

b) The distance from the origin is

$$\frac{8}{5} = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \cdot \underline{r}$$

or simply

$$\begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \cdot \begin{pmatrix} 7 \\ 3 \\ -5 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \cdot \underline{r}$$

a) Using the formula $\underline{r} \cdot \underline{n} = \underline{a} \cdot \underline{n}$ the equation of the plane is

Exercises

1. Find the equation of a plane which is normal to $8\mathbf{i} + 9\mathbf{j} + \mathbf{k}$ and which is a distance 1 from the origin. Give both vector and cartesian forms.
2. Find the equation of a plane which passes through $(8,1,0)$ and which is normal to the vector $\mathbf{i} + 2\mathbf{j} - 3\mathbf{k}$.
3. What is the distance of the plane $\mathbf{r} \cdot \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix} = 5$ from the origin?

Answers

1. $\mathbf{r} \cdot \frac{9\mathbf{i} + \mathbf{j} + 8\mathbf{k}}{\sqrt{146}} = 1$; $8x + 9y + z = \sqrt{146}$.
2. $\mathbf{r} \cdot \begin{pmatrix} 1 \\ 2 \\ -3 \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \\ -3 \end{pmatrix} \cdot \begin{pmatrix} 8 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \\ -3 \end{pmatrix} \cdot \mathbf{r}$, that is $x - 2y + 3z = 10$.
3. $\frac{5\sqrt{14}}{17}$.