

Exponential and Logarithmic Functions



HELM: Helping Engineers Learn Mathematics

http://helm.lboro.ac.uk

About the HELM Project

HELM (Helping Engineers Learn Mathematics) materials were the outcome of a three-year curriculum development project undertaken by a consortium of five English universities led by Loughborough University, funded by the Higher Education Funding Council for England under the Fund for the Development of Teaching and Learning for the period October 2002 – September 2005, with additional transferability funding October 2005 – September 2006.

HELM aims to enhance the mathematical education of engineering undergraduates through flexible learning resources, mainly these Workbooks.

HELM learning resources were produced primarily by teams of writers at six universities: Hull, Loughborough, Manchester, Newcastle, Reading, Sunderland.

HELM gratefully acknowledges the valuable support of colleagues at the following universities and colleges involved in the critical reading, trialling, enhancement and revision of the learning materials:

Aston, Bournemouth & Poole College, Cambridge, City, Glamorgan, Glasgow, Glasgow Caledonian, Glenrothes Institute of Applied Technology, Harper Adams, Hertfordshire, Leicester, Liverpool, London Metropolitan, Moray College, Northumbria, Nottingham, Nottingham Trent, Oxford Brookes, Plymouth, Portsmouth, Queens Belfast, Robert Gordon, Royal Forest of Dean College, Salford, Sligo Institute of Technology, Southampton, Southampton Institute, Surrey, Teesside, Ulster, University of Wales Institute Cardiff, West Kingsway College (London), West Notts College.

HELM Contacts:

Post: HELM, Mathematics Education Centre, Loughborough University, Loughborough, LE11 3TU. *Email:* helm@lboro.ac.uk *Web:* http://helm.lboro.ac.uk

HELM Workbooks List

1	Basic Algebra	26	Functions of a Complex Variable
2	Basic Functions	27	Multiple Integration
3	Equations, Inequalities & Partial Fractions	28	Differential Vector Calculus
4	Trigonometry	29	Integral Vector Calculus
5	Functions and Modelling	30	Introduction to Numerical Methods
6	Exponential and Logarithmic Functions	31	Numerical Methods of Approximation
7	Matrices	32	Numerical Initial Value Problems
8	Matrix Solution of Equations	33	Numerical Boundary Value Problems
9	Vectors	34	Modelling Motion
10	Complex Numbers	35	Sets and Probability
11	Differentiation	36	Descriptive Statistics
12	Applications of Differentiation	37	Discrete Probability Distributions
13	Integration	38	Continuous Probability Distributions
14	Applications of Integration 1	39	The Normal Distribution
15	Applications of Integration 2	40	Sampling Distributions and Estimation
16	Sequences and Series	41	Hypothesis Testing
17	Conics and Polar Coordinates	42	Goodness of Fit and Contingency Tables
18	Functions of Several Variables	43	Regression and Correlation
19	Differential Equations	44	Analysis of Variance
20	Laplace Transforms	45	Non-parametric Statistics
21	z-Transforms	46	Reliability and Quality Control
22	Eigenvalues and Eigenvectors	47	Mathematics and Physics Miscellany
23	Fourier Series	48	Engineering Case Study
24	Fourier Transforms	49	Student's Guide
25	Partial Differential Equations	50	Tutor's Guide

© Copyright Loughborough University, 2015

Production of this 2015 edition, containing corrections and minor revisions of the 2008 edition, was funded by the **sigma** Network.





Exponential and Logarithmic Functions

6.1	The Exponential Function	2
6.2	The Hyperbolic Functions	11
6.3	Logarithms	19
6.4	The Logarithmic Function	27
6.5	Modelling Exercises	38
6.6	Log-linear Graphs	58

Learning outcomes

In this Workbook you will learn about one of the most important functions in mathematics, science and engineering - the exponential function. You will learn how to combine exponential functions to produce other important functions, the hyperbolic functions, which are related to the trigonometric functions.

You will also learn about logarithms and the logarithmic function which is the function inverse to the exponential function. Finally you will learn what a log-linear graph is and how it can be used to simplify the presentation of certain kinds of data.

The Exponential Function





Introduction

In this Section we revisit the use of exponents. We consider how the expression a^x is defined when a is a positive number and x is *irrational*. Previously we have only considered examples in which x is a *rational* number. We consider these exponential functions $f(x) = a^x$ in more depth and in particular consider the special case when the base a is the exponential constant e where :

e = 2.7182818...

We then examine the behaviour of e^x as $x \to \infty$, called **exponential growth** and of e^{-x} as $x \to \infty$ called **exponential decay**.

Prerequisites	 have a good knowledge of indices and their laws 	
Before starting this Section you should	 have knowledge of rational and irrational numbers 	
	• approximate a^x when x is irrational	
Learning Outcomes	 describe the behaviour of a^x: in particular the exponential function e^x 	
On completion you should be able to	 understand the terms exponential growth and exponential decay 	,



1. Exponents revisited

We have seen (in HELM 1.2) the meaning to be assigned to the expression a^p where a is a positive number. We remind the reader that 'a' is called the **base** and 'p' is called the **exponent** (or power or index). There are various cases to consider:

If m, n are positive integers

- $a^n = a \times a \times \cdots \times a$ with n terms
- $a^{1/n}$ means the n^{th} root of a. That is, $a^{1/n}$ is that positive number which satisfies

$$(a^{1/n}) \times (a^{1/n}) \times \dots \times (a^{1/n}) = a$$

where there are \boldsymbol{n} terms on the left hand side.

- $a^{m/n} = (a^{1/n}) \times (a^{1/n}) \times \cdots \times (a^{1/n})$ where there are m terms.
- $a^{-n} = \frac{1}{a^n}$

For convenience we again list the basic laws of exponents:





Solution

First we simplify the numerator:

$$p^{n-2}p^m = p^{n-2+m}$$

Next we simplify the denominator:

$$p^3 p^{2m} = p^{3+2m}$$

Now we combine them and simplify:

$$\frac{p^{n-2}p^m}{p^3p^{2m}} = \frac{p^{n-2+m}}{p^{3+2m}} = p^{n-2+m}p^{-3-2m} = p^{n-2+m-3-2m} = p^{n-m-5}$$



Simplify the expression $\ \ \frac{b^{m-n}b^3}{b^{2m}}$

First simplify the numerator:

Your solution

 $b^{m-n}b^3 =$

Answer

 $b^{m-n}b^3 = b^{m+3-n}$

Now include the denominator:

Your solution
$$\frac{b^{m-n}b^3}{b^{2m}} = \frac{b^{m+3-n}}{b^{2m}} =$$

Answer

 b^{m+3-n} $\frac{b^{2m}}{b^{2m}} = b^{m+3-n-2m} = b^{3-m-n}$



Simplify the numerator:

Your solution $(5a^m)^2 a^2 =$

Answer

$$(5a^m)^2a^2 = 25a^{2m}a^2 = 25a^{2m+2}$$

Now include the denominator:

$$\frac{(5a^m)^2a^2}{(a^3)^2} = \frac{25a^{2m+2}}{a^6} =$$

Answer $\frac{(5a^m)^2a^2}{(a^3)^2} = \frac{25a^{2m+2}}{a^6} = 25a^{2m+2-6} = 25a^{2m-4}$



a^x when x is any real number

So far we have given the meaning of a^p where p is an integer or a rational number, that is, one which can be written as a quotient of integers. So, if p is rational, then

$$p = \frac{m}{n}$$
 where m, n are integers

Now consider x as a real number which cannot be written as a rational number. Two common examples of these **irrational** numbers are $\sqrt{2}$ and π . What we shall do is *approximate* x by a rational number by working to a fixed number of decimal places. For example if

$$x = 3.14159265...$$

then, if we are working to 3 d.p. we would write

$$x \approx 3.142$$

and this number can certainly be expressed as a rational number:

$$x \approx 3.142 = \frac{3142}{1000}$$

so, in this case

 $a^x = a^{3.14159...} \approx a^{3.142} = a^{\frac{3142}{1000}}$

and the final term: $a^{\frac{3142}{1000}}$ can be determined in the usual way by calculator. Henceforth we shall therefore assume that the expression a^x is defined for all positive values of a and for all real values of x.



By working to 3 d.p. find, using your calculator, the value of $3^{\pi/2}$.

First, approximate the value of $\frac{\pi}{2}$:

Your solution $\frac{\pi}{2} \approx$	to 3 d.p.		
$\boxed{ \begin{array}{c} \textbf{Answer} \\ \frac{\pi}{2} \approx \frac{3.1415927\ldots}{2} \end{array} }$	$= 1.5707963 \cdots \approx 1.571$		
Now determine $3^{\pi/2}$: Your solution $3^{\pi/2} \approx$			

Answer

 $3^{\pi/2} \approx 3^{1.571} = 5.618$ to 3 d.p.

2. Exponential functions

For a fixed value of the base a the expression a^x clearly varies with the value of x: it is a function of x. We show in Figure 1 the graphs of $(0.5)^x$, $(0.3)^x$, 1^x , 2^x and 3^x .

The functions a^x (as different values are chosen for a) are called **exponential functions**. From the graphs we see (and these are true for *all* exponential functions):

If
$$a > b > 0$$
 then



Figure 1: $y = a^x$ for various values of a

The most important and widely used exponential function has the particular base e = 2.7182818...It will not be clear to the reader why this particular value is so important. However, its importance will become clear as your knowledge of mathematics increases. The number e is as important as the number π and, like π , is also irrational. The approximate value of e is stored in most calculators. There are numerous ways of calculating the value of e. For example, it can be shown that the value of e is the end-point of the sequence of numbers:

$$\left(\frac{2}{1}\right)^1$$
, $\left(\frac{3}{2}\right)^2$, $\left(\frac{4}{3}\right)^3$, ..., $\left(\frac{17}{16}\right)^{16}$, ..., $\left(\frac{65}{64}\right)^{64}$,...

which, in decimal form (each to 6 d.p.) are

 $2.000000, \quad 2.250000, \quad 2.370370, \quad \dots, \quad 2.637929, \quad \dots, 2.697345, \quad \dots$

This is a slowly converging sequence. However, it does lead to a precise definition for the value of e:

$$\mathbf{e} = \lim_{n \to \infty} \left(\frac{n+1}{n} \right)^n$$

 \mathcal{X}



An quicker way of calculating e is to use the (infinite) series:

$$\mathbf{e} = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \dots + \frac{1}{n!} + \dots$$

where, we remember,

 $n! = n \times (n-1) \times (n-2) \times \dots (3) \times (2) \times (1)$

(This is discussed more fully in HELM 16: Sequences and Series.)

Although all functions of the form a^x are called exponential functions we usually refer to e^x as *the* exponential function.







Exponential functions (and variants) appear in various areas of mathematics and engineering. For example, the shape of a hanging chain or rope, under the effect of gravity, is well described by a combination of the exponential curves e^{kx} , e^{-kx} . The function e^{-x^2} plays a major role in statistics; it being fundamental in the important normal distribution which describes the variability in many naturally occurring phenomena. The exponential function e^{-kx} appears directly, again in the area of statistics, in the Poisson distribution which (amongst other things) is used to predict the number of events (which occur randomly) in a given time interval.

From now on, when we refer to an exponential function, it will be to the function e^x (Figure 2) that we mean, unless stated otherwise.



Use a calculator to determine the following values correct to 2 d.p. (a) $e^{1.5}$, (b) e^{-2} , (c) e^{17} .

Your solution (a) $e^{1.5} =$	(b) $e^{-2} =$	(c) $e^{17} =$	
Answer (a) $e^{1.5} = 4.48$,	(b) $e^{-2} = 0.14$, (c) $e^{17} = 2.4 \times 10^{-2}$	07	



Simplify the expression $\frac{e^{2.7}e^{-3(1.2)}}{e^2}$ and determine its numerical value to 3 d.p.

First simplify the expression:

Your solution $\frac{e^{2.7}e^{-3(1.2)}}{e^2} =$

 $\frac{\mathsf{Answer}}{\mathsf{e}^{2.7}\mathsf{e}^{-3(1.2)}}{\mathsf{e}^2} = \mathsf{e}^{2.7}\mathsf{e}^{-3.6}\mathsf{e}^{-2} = \mathsf{e}^{2.7-3.6-2} = \mathsf{e}^{-2.9}$

Now evaluate its value to 3 d.p.:

Your solution $e^{-2.9} =$ Answer 0.055

3. Exponential growth

If a > 1 then it can be shown that, no matter how large K is:

$$\frac{a^x}{x^K} \to \infty \quad \text{as} \quad x \to \infty$$

That is, if K is fixed (though chosen as large as desired) then eventually, as x increases, a^x will become larger than the value x^K provided a > 1. The growth of a^x as x increases is called **exponential growth**.





A function f(x) grows exponentially and is such that f(0) = 1 and f(2) = 4. Find the exponential curve that fits through these points. Assume the function is $f(x) = e^{kx}$ where k is to be determined from the given information. Find the value of k.

First, find f(0) and f(2) by substituting in $f(x) = e^{kx}$:

Your solution			
When $x = 0$ $f(0) = e^{0}$	0 = 1		
When $x = 2$, $f(2) = 4$	so $e^{2k} = 4$		
By trying values of $k: 0.6$	$6, 0.7, 0.8, {\sf find \ the \ value \ su}$	$ \text{ich that } e^{2k} \approx 4: $	
Your solution			
$e^{2(0.6)} =$	$e^{2(0.7)} =$	$e^{2(0.8)} =$	
Answer			
$e^{2(0.6)} = 3.32$ (too low)	$e^{2(0.7)} = 4.055$ (too hig	gh)	
Now try values of k : $k =$	0.67, 0.68, 0.69:		
Your solution			
$e^{2(0.67)} =$	$e^{2(0.68)} =$	$e^{2(0.69)} =$	
Answer			
$e^{2(0.67)} = 3.819$ (low)	$e^{2(0.68)} = 3.896$ (low)	$e^{2(0.69)} = 3.975$ (low)	
Next try values of $k = 0.6$	691, 0.692:		
Your solution			
$e^{2(0.691)} =$	$e^{2(0.692)} =$	$e^{2(0.693)} =$	
Answer			
$e^{2(0.691)} = 3.983$, (low)	$e^{2(0.692)} = 3.991 \text{ (low)}$	$e^{2(0.693)} = 3.999$ (low)	
Finally, state the exponen	tial function with k to 3 d	.p. which most closely satisfies th	e conditions:
Your solution			

y =

Answer

The exponential function is $e^{0.693x}$.

We shall meet, in Section 6.4, a much more efficient way of finding the value of k.

4. Exponential decay

As we have noted, the behaviour of e^x as $x \to \infty$ is called exponential growth. In a similar manner we characterise the behaviour of the function e^{-x} as $x \to \infty$ as **exponential decay**. The graphs of e^x and e^{-x} are shown in Figure 3.



Figure 3: $y = e^x$ and $y = e^{-x}$

Exponential growth is very rapid and similarly exponential decay is also very rapid. In fact e^{-x} tends to zero so quickly as $x \to \infty$ that, no matter how large the constant K is,

 $x^{K} \mathrm{e}^{-x} \to 0 \quad \text{as} \quad x \to \infty$

The next Task investigates this.



Choose K = 10 in the expression $x^{K}e^{-x}$ and calculate $x^{K}e^{-x}$ using your calculator for x = 5, 10, 15, 20, 25, 30, 35.

Your sol	ution						
x	5	10	15	20	25	30	35
$x^{10}e^{-x}$							
Answer							
Answer x	5	10	15	20	25	30	35

The topics of exponential growth and decay are explored further in Section 6.5.

Exercises

1. Find, to 3 d.p., the values of

(a)
$$2^{-8}$$
 (b) $(5.1)^4$ (c) $(2/10)^{-3}$ (d) $(0.111)^6$ (e) $2^{1/2}$ (f) π^{π} (g) $e^{\pi/4}$ (h) $(1.71)^{-1.71}$

2. Plot $y = x^3$ and $y = e^x$ for 0 < x < 7. For which integer values of x is $e^x > x^3$?

Answers

1. (a) 0.004 (b) 676.520 (c) 125 (d) 0.0 (e) 1.414 (f) 36.462 (g) 2.193 (h) 0.400 2. For integer values of x, $e^x > x^3$ if $x \ge 5$

The Hyperbolic Functions





Introduction

The hyperbolic functions $\sinh x$, $\cosh x$, $\tanh x$ etc are certain combinations of the exponential functions e^x and e^{-x} . The notation implies a close relationship between these functions and the trigonometric functions $\sin x$, $\cos x$, $\tan x$ etc. The close relationship is algebraic rather than geometrical. For example, the functions $\cosh x$ and $\sinh x$ satisfy the relation

 $\cosh^2 x - \sinh^2 x \equiv 1$

which is very similar to the trigonometric identity $\cos^2 x + \sin^2 x \equiv 1$. (In fact every trigonometric identity has an equivalent hyperbolic function identity.)

The hyperbolic functions are not introduced because they are a mathematical nicety. They arise naturally and sufficiently often to warrant sustained study. For example, the shape of a chain hanging under gravity is well described by \cosh and the deformation of uniform beams can be expressed in terms of \tanh .

	 have a good knowledge of the exponential function 	
Prerequisites	• have knowledge of odd and even functions	
Before starting this Section you should	• have familiarity with the definitions of tan, sec, cosec, cot and of trigonometric identities	
	• explain how hyperbolic functions are defined in terms of exponential functions	
Learning Outcomes	• obtain and use hyperbolic function identities	
On completion you should be able to	 manipulate expressions involving hyperbolic functions 	,

1. Even and odd functions

Constructing even and odd functions

A given function f(x) can always be split into two parts, one of which is even and one of which is odd. To do this write f(x) as $\frac{1}{2}[f(x) + f(x)]$ and then simply add and subtract $\frac{1}{2}f(-x)$ to this to give

$$f(x) = \frac{1}{2}[f(x) + f(-x)] + \frac{1}{2}[f(x) - f(-x)]$$

The term $\frac{1}{2}[f(x) + f(-x)]$ is **even** because when x is replaced by -x we have $\frac{1}{2}[f(-x) + f(x)]$ which is the same as the original. However, the term $\frac{1}{2}[f(x) - f(-x)]$ is **odd** since, on replacing x by -x we have $\frac{1}{2}[f(-x) - f(x)] = -\frac{1}{2}[f(x) - f(-x)]$ which is the negative of the original.



Solution $f(x) = x^{3} + 2^{x}$ $f(-x) = (-x)^{3} + 2^{-x} = -x^{3} + 2^{-x}$ Even part: $\frac{1}{2}(f(x) + f(-x)) = \frac{1}{2}(x^{3} + 2^{x} - x^{3} + 2^{-x}) = \frac{1}{2}(2^{x} + 2^{-x})$ Odd part: $\frac{1}{2}(f(x) - f(-x)) = \frac{1}{2}(x^{3} + 2^{x} + x^{3} - 2^{-x}) = \frac{1}{2}(2x^{3} + 2^{x} - 2^{-x})$



Separate the function $x^2 - 3^x$ into odd and even parts.

First, define f(x) and find f(-x):

Your solution		
f(x) =	f(-x) =	
Answer		
$f(x) = x^2 - 3^x$, $f(-x) =$	$x^2 - 3^{-x}$	



Now construct $\frac{1}{2}[f(x) + f(-x)], \frac{1}{2}[f(x) - f(-x)]:$

Your solution

$$\frac{1}{2}[f(x) + f(-x)] = \frac{1}{2}[f(x) - f(-x)] =$$

Answer

$$\begin{split} \frac{1}{2}[f(x) + f(-x)] &= \frac{1}{2}(x^2 - 3^x + x^2 - 3^{-x}) \\ &= x^2 - \frac{1}{2}(3^x + 3^{-x}). \text{ This is the even part of } f(x), \\ \frac{1}{2}[f(x) - f(-x)] &= \frac{1}{2}(x^2 - 3^x - x^2 + 3^{-x}) \\ &= \frac{1}{2}(3^{-x} - 3^x). \text{ This is the odd part of } f(x). \end{split}$$

The odd and even parts of the exponential function

Using the approach outlined above we see that the even part of e^x is

$$\frac{1}{2}(\mathsf{e}^x + \mathsf{e}^{-x})$$

and the odd part of e^x is

$$\frac{1}{2}(\mathsf{e}^x-\mathsf{e}^{-x})$$

We give these new functions special names: $\cosh x$ (pronounced 'cosh' x) and $\sinh x$ (pronounced 'shine' x).



These two functions, when added and subtracted, give

 $\cosh x + \sinh x \equiv e^x$ and $\cosh x - \sinh x \equiv e^{-x}$

The graphs of $\cosh x$ and $\sinh x$ are shown in Figure 4.



Figure 4: $\sinh x$ and $\cosh x$

Note that $\cosh x > 0$ for all values of x and that $\sinh x$ is zero only when x = 0.

2. Hyperbolic identities

The hyperbolic functions $\cosh x$, $\sinh x$ satisfy similar (but not exactly equivalent) identities to those satisfied by $\cos x$, $\sin x$. We note first some basic notation similar to that employed with trigonometric functions:

 $\cosh^n x$ means $(\cosh x)^n$ $\sinh^n x$ means $(\sinh x)^n$ $n \neq -1$

In the special case that n = -1 we **do not** use $\cosh^{-1} x$ and $\sinh^{-1} x$ to mean $\frac{1}{\cosh x}$ and $\frac{1}{\sinh x}$ respectively. The notation $\cosh^{-1} x$ and $\sinh^{-1} x$ is reserved for the **inverse functions** of $\cosh x$ and $\sinh x$ respectively.

Task Show that
$$\cosh^2 x - \sinh^2 x \equiv 1$$
 for all x .

(a) First, express $\cosh^2 x$ in terms of the exponential functions e^x , e^{-x} :

Your solution

$$\cosh^2 x \equiv \left[\frac{1}{2}(e^x + e^{-x})\right]^2 \equiv$$
Answer

$$\frac{1}{4}(e^x + e^{-x})^2 \equiv \frac{1}{4}[(e^x)^2 + 2e^x e^{-x} + (e^{-x})^2] \equiv \frac{1}{4}[e^{2x} + 2e^{x-x} + e^{-2x}] \equiv \frac{1}{4}[e^{2x} + 2 + e^{-2x}]$$



(b) Similarly, express $\sinh^2 x$ in terms of e^x and e^{-x} :



As an alternative to the calculation in this Task we could, instead, use the relations

 $e^x \equiv \cosh x + \sinh x$ $e^{-x} \equiv \cosh x - \sinh x$

and remembering the algebraic identity $(a+b)(a-b) \equiv a^2 - b^2$, we see that

 $(\cosh x + \sinh x)(\cosh x - \sinh x) \equiv e^x e^{-x} \equiv 1$ that is $\cosh^2 x - \sinh^2 x \equiv 1$



The fundamental identity relating hyperbolic functions is:

$$\cosh^2 x - \sinh^2 x \equiv 1$$

This is the hyperbolic function equivalent of the trigonometric identity: $\cos^2 x + \sin^2 x \equiv 1$



Show that $\cosh(x+y) \equiv \cosh x \cosh y + \sinh x \sinh y$.

First, express $\cosh x \cosh y$ in terms of exponentials:

Your solution

$$\cosh x \cosh y \equiv \left(\frac{e^{x} + e^{-x}}{2}\right) \left(\frac{e^{y} + e^{-y}}{2}\right) \equiv$$

$$\boxed{\begin{array}{l} \textbf{Answer}\\ \left(\frac{e^{x} + e^{-x}}{2}\right) \left(\frac{e^{y} + e^{-y}}{2}\right) \equiv \frac{1}{4} [e^{x}e^{y} + e^{-x}e^{y} + e^{x}e^{-y} + e^{-x}e^{-y}] \equiv \frac{1}{4} (e^{x+y} + e^{-x+y} + e^{x-y} + e^{-x-y}) \end{array}}$$

Now express $\sinh x \sinh y$ in terms of exponentials:

Your solution $\left(\frac{e^x - e^{-x}}{2}\right) \left(\frac{e^y - e^{-y}}{2}\right) \equiv$

Answer

$$\left(\frac{\mathsf{e}^{x}-\mathsf{e}^{-x}}{2}\right)\left(\frac{\mathsf{e}^{y}-\mathsf{e}^{-y}}{2}\right) \equiv \frac{1}{4}(\mathsf{e}^{x+y}-\mathsf{e}^{-x+y}-\mathsf{e}^{x-y}+\mathsf{e}^{-x-y})$$

Now express $\cosh x \cosh y + \sinh x \sinh y$ in terms of a hyperbolic function:

Your solution $\cosh x \cosh y + \sinh x \sinh y =$ Answer $\cosh x \cosh y + \sinh x \sinh y \equiv \frac{1}{2}(e^{x+y} + e^{-(x+y)})$ which we recognise as $\cosh(x+y)$



Other hyperbolic function identities can be found in a similar way. The most commonly used are listed in the following Key Point.



3. Related hyperbolic functions

Given the trigonometric functions $\cos x$, $\sin x$ related functions can be defined; $\tan x$, $\sec x$, $\csc x$ through the relations:

 $\tan x \equiv \frac{\sin x}{\cos x} \qquad \qquad \sec x \equiv \frac{1}{\cos x} \qquad \qquad \csc x \equiv \frac{1}{\sin x} \qquad \qquad \cot x \equiv \frac{\cos x}{\sin x}$

In an analogous way, given $\cosh x$ and $\sinh x$ we can introduce hyperbolic functions $\tanh x$, $\operatorname{sec} h x$, $\operatorname{cosech} x$ and $\coth x$. These functions are defined in the following Key Point:





Use the identity $\cosh^2 x - \sinh^2 x \equiv 1$:

Your solution

Answer

Dividing both sides by $\cosh^2 x$ gives

 $1 - \frac{\sinh^2 x}{\cosh^2 x} \equiv \frac{1}{\cosh^2 x}$ implying (see Key Point 6) $1 - \tanh^2 x \equiv \operatorname{sech}^2 x$

Exercises

1. Express

- (a) $2\sinh x + 3\cosh x$ in terms of e^x and e^{-x} .
- (b) $2\sinh 4x 7\cosh 4x$ in terms of e^{4x} and e^{-4x} .

2. Express

- (a) $2e^x e^{-x}$ in terms of $\sinh x$ and $\cosh x$.
- (b) $\frac{7e^x}{(e^x e^{-x})}$ in terms of $\sinh x$ and $\cosh x$, and then in terms of $\coth x$.
- (c) $4e^{-3x} 3e^{3x}$ in terms of $\sinh 3x$ and $\cosh 3x$.
- 3. Using only the \cosh and \sinh keys on your calculator (or e^x key) find the values of
 - (a) tanh 0.35, (b) cosech 2, (c) sech 0.6.

Answers

1. (a)
$$\frac{5}{2}e^x + \frac{1}{2}e^{-x}$$
 (b) $-\frac{5}{2}e^{4x} - \frac{9}{2}e^{-4x}$
2. (a) $\cosh x + 3\sinh x$, (b) $\frac{7(\cosh x + \sinh x)}{2\sinh x}$, $\frac{7}{2}(\coth x + 1)$ (c) $\cosh 3x - 7\sinh 3x$
3. (a) 0.3364, (b) 0.2757 (c) 0.8436



Logarithms





In this Section we introduce the logarithm: $\log_a b$. The operation of taking a logarithm essentially reverses the operation of raising a number to a power. We will formulate the basic laws satisfied by all logarithms and learn how to manipulate expressions involving logarithms. We shall see that to every law of indices there is an equivalent law of logarithms. Although logarithms to any positive base are defined it is common practice to employ only two kinds of logarithms: logs to base 10 and logs to base e.

Prerequisites

Before starting this Section you should ...

Learning Outcomes

On completion you should be able to \ldots

- have a knowledge of exponents and of the laws of indices
- invert $b = a^n$ using logarithms
- simplify expressions involving logarithms
- change bases in logarithms

1. Logarithms

Logarithms reverse the process of raising a base 'a' to a power 'n'. As with all exponentials, the base should be a positive number.

If $b = a^n$ then we write $\log_a b = n$.

Of course, the reverse statement is equivalent

If $\log_a b = n$ then $b = a^n$

The expression $\log_a b = n$ is read

"The log to base a of the number b is equal to n"

The term "log" is short for the word logarithm.



Solution

- (a) Since $16 = 2^4$ then $\log_2 16 = 4$ (b) Since $16 = 4^2$ then $\log_4 16 = 2$
- (c) Since $1000 = 10^3$ then $\log_{10} 1000 = 3$
- (d) Since $134.896 = 10^{2.13}$ then $\log_{10} 134.896 = 2.13$
- (e) Since $8.41467 = e^{2.13}$ then $\log_e 8.414867 = 2.13$





Find the log equivalent of (a) $100 = 10^2$ (b) $\frac{1}{1000} = 10^{-3}$

Here, on the right-hand sides, the base is 10 in each case so:



 $n + m = \log_a(bc)$

From the last Task we have found, using the property of indices, that

 $\log_a(bc) = n + m = \log_a b + \log_a c.$

We conclude that the index law $a^n a^m = a^{n+m}$ has an equivalent logarithm law

 $\log_a(bc) = \log_a b + \log_a c$

In words: "The log of a product is the sum of logs."

Indeed this property is one of the major advantages of using logarithms. They transform a **product** of numbers (a relatively difficult operation) to a **sum** of numbers (a relatively easy operation). Each index law has an equivalent logarithm law, true for any base, listed in the following Key Point:



2. Simplifying expressions involving logarithms

To simplify an expression involving logarithms their laws, given in Key Point 8, need to be used.







Simplify the expression:

$$\log_{10}(\frac{1}{10}) - \log_{10}(\frac{10}{27}) + \log_{10}1000$$

(a) First simplify $\log_{10}(\frac{1}{10})$:

Your solution

 $\log_{10}(\frac{1}{10}) =$

Answer

 $\log_{10}(\frac{1}{10}) = \log_{10} 1 - \log_{10} 10 = 0 - 1 = -1$

(b) Now simplify $\log_{10}(\frac{10}{27})$:

Your solution $\log_{10}(\frac{10}{27}) =$

Answer

 $\log_{10}(\frac{10}{27}) = \log_{10} 10 - \log_{10} 27 = 1 - \log_{10} 27$

(c) Now simplify $\log_{10} 1000$:

Your solution

Answer

3

(d) Finally collect all the terms together from (a), (b), (c) and simplify:

Your solution

Answer

 $-1 - (1 - \log_{10} 27) + 3 = 1 + \log_{10} 27$

3. Logs to base 10 and natural logs

In practice only two kinds of logarithms are commonly used, those to base 10, written \log_{10} (or just simply \log) and those to base e, written \log_e or more usually \ln (called **natural logarithms**). Most scientific calculators will determine the logarithm to base 10 and to base e. For example,

(implying $10^{1.11394} = 13$), (implying $e^{3.13549} = 23$) $\ln 23 = 3.13549$ $\log 13 = 1.11394$



Use your calculator to determine (a) $\log 10$, (b) $\log 1000000$, (c) $\log 0.1$

Your solution
(a) $\log 10 =$ (b) $\log 1000000 =$ (c) $\log 0.1 =$ Answer
(a) 1, (b) 6, (c) -1.Each of the above results could be determined directly, without the use of a calculator. For example:
Since $\log_a a = 1$ then $\log 10 (\equiv \log_{10} 10) = 1$.
Since $\log_a A^k = k \log_a A$ then $\log 1000000 = \log 10^6 = 6 \log 10 = 6$.
Since $\log_a (\frac{A}{B}) = \log_a A - \log_a B$ and $\log_a 1 = 0$ and $\log_a a = 0$, then
 $\log 0.1 = \log(\frac{1}{10}) = \log 1 - \log(10) = -1$ Use your calculator to determine
(a) $\ln 29.42$, (b) $\ln e$, (c) $\ln 0.1$ Your solution

(a) $\ln 29.42 =$ (b) $\ln e =$ (c) $\ln 0.1 =$

Answer

(a) $\ln 29.42 = 3.38167$, (b) $\ln e = 1$, (c) $\ln 0.1 = -2.30258$

4. Changing base in logarithms

It is sometimes required to express the logarithm with respect to one base in terms of a logarithm with respect to another base.

Now

 $b = a^n$ implies $\log_a b = n$

where we have used logs to base a. What happens if, for some reason, we want to use another base, p say? We take logs (to base p) of both sides of $b = a^n$:

 $\log_p(b) = \log_p(a^n) = n \log_p a$ (using one of the logarithm laws)

So

$$n = \frac{\log_p(b)}{\log_p(a)}$$
 that is $\log_a b = \frac{\log_p(b)}{\log_p(a)}$

This is the rule to be used when converting logarithms from one base to another.





For base 10 logs:

 $\log_a b = \frac{\log(b)}{\log(a)}$

For example,

$$\log_3 7 = \frac{\log 7}{\log 3} = \frac{0.8450980}{0.4771212} = 1.7712437$$

(Check, on your calculator, that $3^{1.7712437} = 7$). For natural logs:

$$\log_a b = \frac{\ln(b)}{\ln(a)}$$

For example,

$$\log_3 7 = \frac{\ln 7}{\ln 3} = \frac{1.9459101}{1.0986123} = 1.7712437$$

Of course, $\log_3 7$ cannot be determined directly on your calculator since logs to base 3 are not available but it can be found using the above method.



Use your calculator to determine the value of $\log_{21}7$ using first base 10 then check using base e.

Re-express $\log_{21}7$ using base 10 then base e:

Your solution

$$\log_{21} 7 = \frac{\log 7}{\log 21} =$$
 $\log_{21} 7 = \frac{\ln 7}{\ln 21} =$

 Answer
 $\log_{21} 7 = \frac{\log 7}{\log 21} = 0.6391511$
 $\log_{21} 7 = \frac{\ln 7}{\ln 21} = 0.6391511$



Solution

Let $y = 10^{\log x}$ then take logs (to base 10) of both sides: $\log y = \log(10^{\log x}) = (\log x) \log 10$ where we have used: $\log A^k = k \log A$. However, since we are using logs to base 10 then $\log 10 = 1$ and so

 $\log y = \log x \quad \text{ implying } \quad y = x$ Therefore, finally we conclude that

 $10^{\log x} = x$

This is an important result true for logarithms of any base. It follows from the basic definition of the logarithm.



Raising to the power and taking logs are **inverse** operations.

Exercises

- 1. Find the values of (a) $\log_2 8$ (b) $\log_{16} 50$ (c) $\ln 28$
- 2. Simplify
 - (a) $\log 1 3 \log 2 + \log 16$.
 - (b) $10 \log x 2 \log x^2$.
 - (c) $\ln(8x-4) \ln(4x-2)$.
 - (d) $\ln 10 \log 7 \ln 7$.

Answers

1. (a) 3 (b) 1.41096 (c) 3.3322 2. (a) $\log 2$, (b) $6 \log x$ or $\log x^6$, (c) $\ln 2$, (d) 0

The Logarithmic Function





In this Section we consider the logarithmic function $y = \log_a x$ and examine its important characteristics. We see that this function is only defined if x is a positive number. We also see that the \log function is the inverse of the exponential function and vice versa. We show, through numerous examples, how equations involving logarithms and exponentials can be solved.

	• have knowledge of inverse functions	
Prerequisites	 have knowledge of the laws of logarithms and of the laws of indices 	
Defore starting this Section you should	• be able to solve quadratic equations	
Learning Outcomes	 explain the relation between the logarithm and the exponential function 	
On completion you should be able to	 solve equations involving exponentials and logarithms 	/

1. The logarithmic function

In Section 6.3 we introduced the operation of taking logarithms which reverses the operation of exponentiation.

If
$$a > 0$$
 and $a \neq 1$ then $x = a^y$ implies $y = \log_a x$

In this Section we consider the log function in more detail. We shall concentrate only on the functions $\log x$ (i.e. to base 10) and $\ln x$ (i.e. to base e). The functions $y = \log x$ and $y = \ln x$ have similar characteristics. We can never choose x as a negative number since 10^y and e^y are each always positive. The graphs of $y = \log x$ and $y = \ln x$ are shown in Figure 5.



Figure 5: Logarithmic and exponential functions

From the graphs we see that both functions are one-to-one so each has an **inverse function** - the inverse function of $\log_a x$ is a^x . Let us do this for logs to base 10.

2. Solving equations involving logarithms and exponentials

To solve equations which involve logarithms or exponentials we need to be aware of the basic laws which govern both of these mathematical concepts. We illustrate by considering some examples.

Example 6
Solve for the variable
$$x$$
: (a) $3 = 10^x$, (b) $10^{x/4} = \log 3$, (c) $\frac{1}{17 - e^x} = 4$

Solution

(a) Here we take logs (to base 10 because of the term 10^x) of both sides to get

$$\log 3 = \log 10^x = x \log 10 = x$$

where we have used the general property that $\log_a A^k = k \log_a A$ and the specific property that $\log 10 = 1$. Hence $x = \log 3$ or, in numerical form, x = 0.47712 to 5 d.p.



Solution (contd.)

(b) The approach used in (a) is used here. Take logs of both sides: $\log(10^{x/4}) = \log(\log 3)$

that is
$$\frac{x}{4}\log 10 = \log(\log 3) = \log(0.4771212) = -0.3213712$$

So, since $\log 10 = 1$, we have x = 4(-0.3213712) = -1.28549 to 5 d.p.

(c) Here we simplify the expression before taking logs.

 $\frac{1}{17 - e^x} = 4$ implies $1 = 4(17 - e^x)$

or $4e^x = 4(17) - 1 = 67$ so $e^x = 16.75$. Now taking natural logs of both sides (because of the presence of the e^x term) we have:

 $\ln(\mathbf{e}^x) = \ln(16.75) = 2.8183983$

But $\ln(e^x) = x \ln e = x$ and so the solution to $\frac{1}{17 - e^x} = 4$ is x = 2.81840 to 5 d.p.



First solve for e^x by taking square roots of both sides:

Your solution $(e^x)^2 = 50$ implies $e^x =$

Answer

 $(e^x)^2 = 50$ implies $e^x = \sqrt{50} = 7.071068$. Here we have taken the positive value for the square root since we know that exponential functions **are always positive**.

Now take logarithms to an appropriate base to find x:

Your solution

 $e^x = 7.071068$ implies x =

Answer

 $e^x = 7.071068$ implies $x = \ln(7.071068) = 1.95601$ to 5 d.p.



Solve the equation $e^{2x} = 17e^x$

First simplify the expression as much as possible (divide both sides by e^x):

Your solution $e^{2x} = 17e^x$ implies $\frac{e^{2x}}{e^x} = 17$ so Answer

 $\frac{e^{2x}}{e^x} = 17$ implies $e^{2x-x} = 17$ so $e^x = 17$

Now complete the solution for x:

Your solution

 $e^x = 17$ implies x =

Answer

 $x = \ln(17) = 2.8332133$



Solution

We first simplify this expression by multiplying through by 10^x (to eliminate the term 10^{-x}):

$$10^{x}(10^{x}) - 10^{x}(5) + 10^{x}(6(10^{-x})) = 0$$

or

 $(10^x)^2 - 5(10^x) + 6 = 0$ since $10^x(10^{-x}) = 10^0 = 1$

We realise that this expression is a quadratic equation. Let us put $y = 10^x$ to give

 $y^2 - 5y + 6 = 0$

Now, we can factorise to give

(y-3)(y-2)=0 so that y=3 or y=2

For each of these values of y we obtain a separate value for x since $y = 10^x$.

Case 1 If y = 3 then $3 = 10^x$ implying $x = \log 3 = 0.4771212$

Case 2 If y = 2 then $2 = 10^x$ implying $x = \log 2 = 0.3010300$

We conclude that the equation $10^x - 5 + 6(10^{-x}) = 0$ has two possible solutions for x: either x = 0.4771212 or x = 0.3010300, to 7 d.p.



Solve
$$2e^{2x} - 7e^x + 3 = 0.$$

First write this equation as a quadratic in the variable $y = e^x$ remembering that $e^{2x} \equiv (e^x)^2$:

Your solution If $y = e^x$ then $2e^{2x} - 7e^x + 3 = 0$ becomes Answer $2y^2 - 7y + 3 = 0$ Now solve the quadratic for *y*: Your solution $2y^2 - 7y + 3 = 0 \quad \text{implies} \quad (2y$)(y) = 0Answer (2y-1)(y-3) = 0 therefore $y = \frac{1}{2}$ or y = 3Finally, for each of your values of y, find x: Your solution If $y = \frac{1}{2}$ then $\frac{1}{2} = e^x$ implies x =If y = 3 then $3 = e^x$ implies x =Answer

x = -0.693147 or x = 1.0986123



The temperature T, in degrees C, of a chemical reaction is given by the formula $T = 80e^{0.03t} \times t \ge 0$, where t is the time, in seconds.

Calculate the time taken for the temperature to reach $150^\circ~{\rm C}$.

Answer $150 = 80e^{0.03t} \Rightarrow 1.875 = e^{0.03t} \Rightarrow \ln(1.875) = 0.03t \Rightarrow t = \frac{\ln(1.875)}{0.03}$ This gives t = 20.95 to 2 d.p. So the time is 21 seconds.



Engineering Example 1

Arrhenius' law

Introduction

Chemical reactions are very sensitive to temperature; normally, the rate of reaction increases as temperature increases. For example, the corrosion of iron and the spoiling of food are more rapid at higher temperatures. Chemically, the probability of collision between two molecules increases with temperature, and an increased collision rate results in higher kinetic energy, thus increasing the proportion of molecules that have the **activation energy** for the reaction, i.e. the minimum energy required for a reaction to occur. Based upon his observations, the Swedish chemist, Svante Arrhenius, proposed that the rate of a chemical reaction increases exponentially with temperature. This relationship, now known as Arrhenius' law, is written as

$$k = k_0 \, \exp\left(\frac{-E_a}{R\,T}\right) \tag{1}$$

where k is the reaction rate constant, k_0 is the frequency factor, E_a is the activation energy, R is the universal gas constant and T is the absolute temperature. Thus, the reaction rate constant, k, depends on the quantities k_0 and E_a , which characterise a given reaction, and are generally assumed to be temperature independent.

Problem in words

In a laboratory, ethyl acetate is reacted with sodium hydroxide to investigate the reaction kinetics. Calculate the frequency factor and activation energy of the reaction from Arrhenius' Law, using the experimental measurements of temperature and reaction rate constant in the table:

T	310	350
k	7.757192	110.9601

Mathematical statement of problem

Given that $k = 7.757192 \text{ s}^{-1}$ at T = 310 K and $k = 110.9601 \text{ s}^{-1}$ at T = 350 K, use Equation (1) to produce two linear equations in E_a and k_0 . Solve these to find E_a and k_0 . (Assume that the gas constant $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$.)

Mathematical analysis

Taking the natural logarithm of both sides of (1)

$$\ln k = \ln \left\{ k_0 \, \exp\left(\frac{-E_a}{RT}\right) \right\} = \ln k_0 - \frac{E_a}{RT}$$

Now inserting the experimental data gives the two linear equations in E_a and k_0

$$\ln k_1 = \ln k_0 - \frac{E_a}{R T_1}$$
(2)

$$\ln k_2 = \ln k_0 - \frac{E_a}{R T_2}$$
(3)

where $k_1 = 7.757192$, $T_1 = 310$ and $k_2 = 110.9601$, $T_2 = 350$.



Firstly, to find E_a , subtract Equation (2) from Equation (3)

$$\ln k_2 - \ln k_1 = \frac{E_a}{RT_1} - \frac{E_a}{RT_2} = \frac{E_a}{R} \left(\frac{1}{T_1} - \frac{1}{T_2}\right)$$

so that

$$E_{a} = \frac{R\left(\ln k_{2} - \ln k_{1}\right)}{\left(\frac{1}{T_{1}} - \frac{1}{T_{2}}\right)}$$

and substituting the values gives

$$E_a = 60000 \text{ J mol}^{-1} = 60 \text{ kJ mol}^{-1}$$

Secondly, to find k_0 , from (2)

$$\ln k_0 = \ln k_1 + \frac{E_a}{RT_1} \qquad \Rightarrow \quad k_0 = \exp\left(\ln k_1 + \frac{E_a}{RT_1}\right) = k_1 \exp\left(\frac{E_a}{RT_1}\right)$$

and substituting the values gives

$$k_0 = 1.0 \times 10^{11} \text{ s}^{-1}$$



The reaction

$$2NO_2(g) \longrightarrow 2NO(g) + O_2(g)$$

has a reaction rate constant of $1.0\times10^{-10}~{\rm s}^{-1}$ at 300 K and activation energy of 111 kJ mol $^{-1}$ = 111 000 J mol $^{-1}$. Use Arrhenius' law to find the reaction rate constant at a temperature of 273 K.

Your solution

Answer Rearranging Arrhenius' equation gives

$$k_0 = k \, \exp\left(\frac{E_a}{R \, T}\right)$$

Substituting the values gives $k_0=2.126\times 10^9~{\rm s}^{-1}$

Now we use this value of k_0 with E_a in Arrhenius' equation (1) to find k at T = 273 K

$$k = k_0 \exp\left(\frac{-E_a}{RT}\right) = 1.226 \times 10^{-12} \text{ s}^{-1}$$



For a chemical reaction with frequency factor $k_0 = 0.5 \text{ s}^{-1}$ and ratio $E_a/R = 800 \text{ K}$, use Arrhenius' law to find the temperature at which the reaction rate constant would be equal to 0.1 s⁻¹.

Your solution

Answer

Rearranging Equation (1)

$$\frac{k}{k_0} = \exp\left(\frac{-E_a}{RT}\right)$$

Taking the natural logarithm of both sides

$$\ln\left(\frac{k}{k_0}\right) = \frac{-E_a}{RT}$$

so that

$$T=\frac{-E_a}{R\,\ln{(k/k_0)}}=\frac{E_a}{R\,\ln{(k_0/k)}}$$
 Substituting the values gives $T=497~{\rm K}$

As a final example we consider equations involving the hyperbolic functions.


(a) $\cosh 3x = 1$ (b) $\cosh 3x = 2$ (c) $2 \cosh^2 x = 3 \cosh 2x - 3$

Solution

- (a) From its graph we know that $\cosh x = 0$ only when x = 0, so we need 3x = 0 which implies x = 0.
- (b) $\cosh 3x = 2$ implies $\frac{e^{3x} + e^{-3x}}{2} = 2$ or $e^{3x} + e^{-3x} 4 = 0$

Now multiply through by e^{3x} (to eliminate the term e^{-3x}) to give

$$e^{3x}e^{3x} + e^{3x}e^{-3x} - 4e^{3x} = 0$$
 or $(e^{3x})^2 - 4e^{3x} + 1 = 0$

This is a quadratic equation in the variable e^{3x} so substituting $y = e^{3x}$ gives

$$y^2 - 4y + 1 = 0$$
 implying $y = 2 \pm \sqrt{3}$ so $y = 3.7321$ or 0.26795
 $e^{3x} = 3.7321$ implies $x = \frac{1}{3} \ln 3.7321 = 0.439$ to 3 d.p.
 $e^{3x} = 0.26795$ implies $x = \frac{1}{3} \ln 0.26795 = -0.439$ to 3 d.p.

(c) We first simplify this expression by using the identity: $\cosh 2x = 2 \cosh^2 -1$. Thus the original equation $2 \cosh^2 x = 3 \cosh 2x - 3$ becomes $\cosh 2x + 1 = 3 \cosh 2x - 3$ or, when written in terms of exponentials:

$$\frac{e^{2x} + e^{-2x}}{2} = 3(\frac{e^{2x} + e^{-2x}}{2}) - 4$$

Multiplying through by $2e^{2x}$ gives $e^{4x} + 1 = 3(e^{4x} + 1) - 8e^{2x}$ or, after simplifying:

$$e^{4x} - 4e^{2x} + 1 = 0$$

Writing $y = e^{2x}$ we easily obtain $y^2 - 4y + 1 = 0$ with solution (using the quadratic formula):

$$y = \frac{4 \pm \sqrt{16 - 4}}{2} = 2 \pm \sqrt{3}$$

If $y = 2 + \sqrt{3}$ then $2 + \sqrt{3} = e^{2x}$ implying $x = 0.65848$ to 5 d.p.
If $y = 2 - \sqrt{3}$ then $2 - \sqrt{3} = e^{2x}$ implying $x = -0.65848$ to 5 d.p.



Find the solution for x if tanh x = 0.5.

First re-write tanh x in terms of exponentials:

Your solution
$\tanh x =$
Answer
Now substitute into $\tanh x = 0.5$:
Your solution $\tanh x = 0.5$ implies $\frac{e^{2x} - 1}{e^{2x} + 1} = 0.5$ so, on simplifying, $e^{2x} =$
Answer $\frac{e^{2x} - 1}{e^{2x} + 1} = 0.5 \text{ implies } (e^{2x} - 1) = \frac{1}{2}(e^{2x} + 1) \text{ so } \frac{e^{2x}}{2} = \frac{3}{2} \text{ so, finally, } e^{2x} = 3$
Now complete your solution by finding <i>x</i> :

Your solution

 $e^{2x} = 3$ so x =

Answer

 $x = \frac{1}{2}\ln 3 = 0.549306$

Alternatively, many calculators can directly calculate the inverse function $tanh^{-1}$. If you have such a calculator then you can use the fact that

 $\tanh x = 0.5$ implies $x = \tanh^{-1} 0.5$ to obtain directly x = 0.549306





Solution

This has logs to two different bases. So we must first express each logarithm in terms of logs to the *same* base, e say. From Key Point 8

$$\log x = \frac{\ln x}{\ln 10}$$

So $3 \ln x + 4 \log x = 1$ becomes
 $3 \ln x + 4 \frac{\ln x}{\ln 10} = 1$ or $(3 + \frac{4}{\ln 10}) \ln x = 1$
leading to $\ln x = \frac{\ln 10}{3 \ln 10 + 4} = \frac{2.302585}{10.907755} = 0.211096$ and so
 $x = e^{0.211096} = 1.2350311$

Exercises

1. Solve for the variable x: (a) $\pi = 10^x$ (b) $10^{-x/2} = 3$ (c) $\frac{1}{17 - \pi^x} = 4$

2. Solve the equations

(a) $e^{2x} = 17e^x$, (b) $e^{2x} - 2e^x - 6 = 0$, (c) $\cosh x = 3$.

Answers

1. (a) $x = \log \pi = 0.497$ (b) $-x/2 = \log 3$ and so $x = -2\log 3 = -0.954$ (c) $17 - \pi^x = 0.25$ so $\pi^x = 16.75$ therefore $x = \frac{\log 16.75}{\log \pi} = \frac{1.224}{0.497} = 2.462$ 2. (a) Take logs of both sides: $2x = \ln 17 + x$ \therefore $x = \ln 17 = 2.833$ (b) Let $y = e^x$ then $y^2 - 2y - 6 = 0$ therefore $y = 1 \pm \sqrt{7}$ (we cannot take the negative sign since exponentials can never be negative). Thus $x = \ln(1 + \sqrt{7}) = 1.2936$. (c) $e^x + e^{-x} = 6$ therefore $e^{2x} - 6e^x + 1 = 0$ so $e^x = \frac{6 \pm \sqrt{36 - 4}}{2} = 3 \pm \sqrt{8}$ We have, finally $x = \ln(3 + \sqrt{8}) = 1.7627$ or $x = \ln(3 - \sqrt{8}) = -1.7627$

Modelling Exercises





This Section provides examples and tasks employing exponential functions and logarithmic functions, such as growth and decay models which are important throughout science and engineering.



1. Exponential increase



- (a) Look back at Section 6.2 to review the definitions of an exponential function and the exponential function.
- (b) List examples in this Workbook of contexts in which exponential functions are appropriate.

Your solution

Answer

- (a) An exponential function has the form $y = a^x$ where a > 0. The exponential function has the form $y = e^x$ where e = 2.718282...
- (b) It is stated that exponential functions are useful when modelling the shape of a hanging chain or rope under the effect of gravity or for modelling exponential growth or decay.

We will look at a specific example of the exponential function used to model a population increase.



Given that

 $P = 12e^{0.1t}$ $(0 \le t \le 25)$

where P is the number in the population of a city **in millions** at time t **in years** answer these questions.

- (a) What does this model imply about P when t = 0?
- (b) What is the stated upper limit of validity of the model?
- (c) What does the model imply about values of P over time?
- (d) What does the model predict for P when t = 10? Comment on this.
- (d) What does the model predict for P when t = 25? Comment on this.

Your	solution
(a)	
(b)	
(c)	
(d)	
(e)	
Ansv	ver
(a)	At $t = 0$, $P = 12$ which represents the initial population of 12 million. (Recall that $e^0 = 1$.)
(b)	The time interval during which the model is valid is stated as $(0 \le t \le 25)$ so the model is intended to apply for 25 years.
(c)	This is exponential growth so P will increase from 12 million at an accelerating rate.
(d)	$P(10)=12{\rm e}^1\approx$ 33 million. This is getting very large for a city but might be attainable in 10 years and just about sustainable.
(e)	$P(25) = 12e^{2.5} \approx 146$ million. This is unrealistic for a city.

Note that exponential population growth of the form $P = P_0 e^{kt}$ means that as t becomes large and positive, P becomes very large. Normally such a population model would be used to predict values of P for t > 0, where t = 0 represents the present or some fixed time when the population is known. In Figure 6, values of P are shown for t < 0. These correspond to extrapolation of the model into the past. Note that as t becomes increasingly negative, P becomes very small but is never zero or negative because e^{kt} is positive for all values of t. The parameter k is called the **instantaneous fractional growth rate**.



Figure 6: The function $P = 12e^{0.01t}$



For the model $P = 12e^{kt}$ we see that k = 0.1 is unrealistic, and more realistic values would be k = 0.01 or k = 0.02. These would be similar but k=0.02 implies a faster growth for t > 0 than k = 0.01. This is clear in the graphs for k = 0.01 and k = 0.02 in Figure 7. The functions are plotted up to 200 years to emphasize the increasing difference as t increases.



Figure 7: Comparison of the functions $P = 12e^{0.01t}$ and $P = 12e^{0.02t}$

The exponential function may be used in models for other types of growth as well as population growth. A general form may be written

$$y = ae^{bx} \qquad a > 0, \quad b > 0, \quad c \le x \le d$$

where a represents the value of y at x = 0. The value a is the intercept on the y-axis of a graphical representation of the function. The value b controls the rate of growth and c and d represent limits on x.

In the general form, a and b represent the **parameters** of the exponential function which can be selected to fit any given modelling situation where an exponential function is appropriate.

2. Linearisation of exponential functions

This subsection relates to the description of log-linear plots covered in Section 6.6.

Frequently in engineering, the question arises of how the parameters of an exponential function might be found from given data. The method follows from the fact that it is possible to 'undo' the exponential function and obtain a linear function by means of the logarithmic function. Before showing the implications of this method, it may be necessary to remind you of some rules for manipulating logarithms and exponentials. These are summarised in Table 1 on the next page, which exactly matches the general list provided in Key Point 8 in Section 6.3 (page 22.)

Number	Rule	Number	Rule
1a	$\ln(xy) = \ln(x) + \ln(y)$	1b	$e^x \times e^y = e^{x+y}$
2a	$\ln(x/y) = \ln(x) - \ln(y)$	2b	$e^x/e^y=e^{x-y}$
3a	$\ln(x^y) = y\ln(x)$	3b	$(e^x)^y = e^{xy}$
4a	$\ln(\mathbf{e}^x) = x$	4b	$e^{\ln(x)} = x$
5a	$\ln(e) = 1$	5b	$e^1 = e$
ба	$\ln(1) = 0$	6b	$e^0 = 1$

Table 1: Rules for manipulating base e logarithms and exponentials

We will try 'undoing' the exponential in the particular example

$$P = 12e^{0.1t}$$

We take the natural logarithm (ln) of both sides, which means logarithm to the base e. So

 $\ln(P) = \ln(12e^{0.1t})$

The result of using Rule 1a in Table 1 is

 $\ln(P) = \ln(12) + \ln(e^{0.1t}).$

The natural logarithmic functions 'undoes' the exponential function, so by Rule 4a,

$$\ln(e^{0.1t}) = 0.1t$$

and the original equation for \boldsymbol{P} becomes

$$\ln(P) = \ln(12) + 0.1t.$$

Compare this with the general form of a linear function y = ax + b.

$$y = ax + b$$

$$\downarrow \qquad \downarrow \qquad \downarrow$$

$$\ln(P) = 0.1t + \ln(12)$$

If we regard $\ln(P)$ as equivalent to y, 0.1 as equivalent to the constant a, t as equivalent to x, and $\ln(12)$ as equivalent to the constant b, then we can identify a linear relationship between $\ln(P)$ and t. A plot of $\ln(P)$ against t should result in a straight line, of slope 0.1, which crosses the $\ln(P)$ axis at $\ln(12)$. (Such a plot is called a **log-linear** or **log-lin** plot.) This is not particularly interesting here because we know the values 12 and 0.1 already.

Suppose, though, we want to try using the general form of the exponential function

 $P = ae^{bt} \qquad (c \le t \le d)$

to create a continuous model for a population for which we have some discrete data. The first thing to do is to take logarithms of both sides

$$\ln(P) = \ln(ae^{bt}) \qquad (c \le t \le d).$$

Rule 1 from Table 1 then gives

$$\ln(P) = \ln(a) + \ln(e^{bt}) \qquad (c \le t \le d).$$

But, by Rule 4a, $\ln(e^{bt}) = bt$, so this means that

$$\ln(P) = \ln(a) + bt \qquad (c \le t \le d).$$



So, given some 'population versus time' data, for which you believe can be modelled by some version of the exponential function, plot the natural logarithm of population against time. If the exponential function is appropriate, the resulting data points should lie on or near a straight line. The slope of the straight line will give an estimate for b and the intercept with the $\ln(P)$ axis will give an estimate for $\ln(a)$. You will have carried out a **logarithmic transformation** of the original data for P. We say the original variation has been **linearised**.

A similar procedure will work also if any exponential function rather than the base e exponential function is used. For example, suppose that we try to use the function

$$P = A \times 2^{Bt} \qquad (C \le t \le D),$$

where A and B are constant parameters to be derived from the given data. We can take natural logarithms again to give

$$\ln(P) = \ln(A \times 2^{Bt}) \qquad (C \le t \le D).$$

Rule 1a from Table 1 then gives

$$\ln(P) = \ln(A) + \ln(2^{Bt}) \qquad (C \le t \le D).$$

Rule 3a then gives

$$\ln(2^{Bt}) = Bt\ln(2) = B\ln(2) t$$

and so

$$\ln(P) = \ln(A) + B\ln(2) t \qquad (C \le t \le D)$$

Again we have a straight line graph with the same intercept as before, $\ln A$, but this time with slope $B \ln(2)$.



The amount of money $\pounds M$ to which $\pounds 1$ grows after earning interest of 5% p.a. for N years is worked out as

 $M = 1.05^{N}$

Find a linearised form of this equation.

Your solution

Answer

Take natural logarithms of both sides.

 $\ln(M) = \ln(1.05^N).$

Rule 3b gives

 $\ln(M) = N\ln(1.05).$

So a plot of $\ln(M)$ against N would be a straight line passing through (0,0) with slope $\ln(1.05)$.

The linearisation procedure also works if logarithms other than natural logarithms are used. We start again with

 $P = A \times 2^{Bt} \qquad (C \le t \le D).$

and will take logarithms to base 10 instead of natural logarithms. Table 2 presents the laws of logarithms and indices (based on Key Point 8 page 22) interpreted for \log_{10} .

Number	Rule	Number	Rule
1a	$\log_{10}(AB) = \log_{10} A + \log_{10} B$	1b	$10^A 10^B = 10^{A+B}$
2a	$\log_{10}(A/B) = \log_{10}A - \log_{10}B$	2b	$10^A / 10^B = 10^{A-B}$
3a	$\log_{10}(A^k) = k \log_{10} A$	3b	$(10^A)^k = 10^{kA}$
4a	$\log_{10}(10^A) = A$	4b	$10^{\log_{10} A} = A$
5a	$\log_{10} 10 = 1$	5b	$10^1 = 10$
ба	$\log_{10} 1 = 0$	6b	$10^0 = 1$

Table 2: Rules for manipulating base 10 logarithms and exponentials

Taking logs of $P = A \times 2^{Bt}$ gives:

$$\log_{10}(P) = \log_{10}(A \times 2^{Bt}) \qquad (C \le t \le D).$$

Rule 1a from Table 2 then gives

 $\log_{10}(P) = \log_{10}(A) + \log_{10}(2^{Bt}) \qquad (C \le t \le D).$

Use of Rule 3a gives the result

 $\log_{10}(P) = \log_{10}(A) + B \log_{10}(2) t \qquad (C \le t \le D).$



(a) Write down the straight line function corresponding to taking logarithms of the general exponential function

$$P = ae^{bt} \qquad (c \le t \le d)$$

by taking logarithms to base 10.

(b) Write down the slope of this line.

Your solution

Answer

(a)
$$\log_{10}(P) = \log_{10}(a) + (b \log_{10}(e))t$$
 $(c \le t \le d)$

(b) $b \log_{10}(e)$

It is not usually necessary to declare the subscript 10 when indicating logarithms to base 10. If you meet the term 'log' it will probably imply "to the base 10". In the remainder of this Section, the subscript 10 is dropped where \log_{10} is implied.



3. Exponential decrease

Consider the value, $\pounds D$, of a car subject to depreciation, in terms of the age A years of the car. The car was bought for $\pounds 10500$. The function

 $D = 10500e^{-0.25A} \qquad (0 \le A \le 6)$

could be considered appropriate on the ground that (a) D had a fixed value of $\pounds 10500$ when A = 0, (b) D decreases as A increases and (c) D decreases faster when A is small than when A is large. A plot of this function is shown in Figure 8.



Figure 8: Plot of car depreciation over 6 years







Engineering Example 2

Exponential decay of sound intensity

Introduction

The rate at which a quantity decays is important in many branches of engineering and science. A particular example of this is exponential decay. Ideally the sound level in a room where there are substantial contributions from reflections at the walls, floor and ceiling will decay exponentially once the source of sound is stopped. The decay in the sound intensity is due to absorbtion of sound at the room surfaces and air absorption although the latter is significant only when the room is very large. The contributions from reflection are known as **reverberation**. A measurement of reverberation in a room of known volume and surface area can be used to indicate the amount of absorption.

Problem in words

As part of an emergency test of the acoustics of a concert hall during an orchestral rehearsal, consultants asked the principal trombone to play a single note at maximum volume. Once the sound had reached its maximum intensity the player stopped and the sound intensity was measured for the next 0.2 seconds at regular intervals of 0.02 seconds. The initial maximum intensity at time 0 was 1. The readings were as follows:

time	0	0.02	0.04	0.06	0.08	0.10	0.12	0.14	0.16	0.18	0.20
intensity	1	0.63	0.35	0.22	0.13	0.08	0.05	0.03	0.02	0.01	0.005

Draw a graph of intensity against time and, assuming that the relationship is exponential, find a function which expresses the relationship between intensity and time.

Mathematical statement of problem

If the relationship is exponential then it will be a function of the form

$$I = I_0 10^{kt}$$

and a log-linear graph of the values should lie on a straight line. Therefore we can plot the values and find the gradient and the intercept of the resulting straight-line graph in order to find the values for I_0 and k.

 \boldsymbol{k} is the gradient of the log-linear graph i.e.

$$k = \frac{\text{change in } \log_{10} \text{ (intensity)}}{\text{change in time}}$$

and I_0 is found from where the graph crosses the vertical axis $\log_{10}(I_0) = c$

Mathematical analysis

Figure 9(a) shows the graph of intensity against time.



We calculate the \log_{10} (intensity) to create the table below:

time	0	0.02	0.04	0.06	0.08	0.10	0.12	0.14	0.16	0.18	0.20
$\log_{10}(\text{intensity})$	0	-0.22	-0.46	-0.66	-0.89	-1.1	-1.3	-1.5	-1.7	-2.0	-2.2

Figure 9(b) shows the graph of \log (intensity) against time.



Figure 9: (a) Graph of sound intensity against time (b) Graph of \log_{10} (intensity) against time and a line fitted by eye to the data. The line goes through the points (0,0) and (0.2, -2.2).

We can see that the second graph is approximately a straight line and therefore we can assume that the relationship between the intensity and time is exponential and can be expressed as

$$I = I_0 10^{kt}.$$

The log_{10} of this gives

 $\log_{10}(I) = \log_{10}(I_0) + kt.$

From the graph (b) we can measure the gradient, k using

$$k = \frac{\text{change in } \log_{10} (\text{intensity})}{\text{change in time}}$$
 giving $k = \frac{-2.2 - 0}{0.2 - 0} = -11$

The point at which it crosses the vertical axis gives

 $\log_{10}(I_0) = 0 \Rightarrow I_0 = 10^0 = 1$

Therefore the expression $I = I_0 10^{kt}$ becomes

$$I = 10^{-11}$$

Interpretation

The data recorded for the sound intensity fit exponential decaying with time. We have used a log-linear plot to obtain the approximate function:

 $I = 10^{-11t}$

4. Growth and decay to a limit

Consider a function intended to represent the speed of a parachutist after the opening of the parachute where $v \text{ m s}^{-1}$ is the instantanous speed at time t s. An appropriate function is

 $v = 12 - 8e^{-1.25t} \qquad (t \ge 0),$

We will look at some of the properties and modelling implications of this function. Consider first the value of v when t = 0:

 $v = 12 - 8e^0 = 12 - 8 = 4$

This means the function predicts that the parachutist is moving at 4 m s⁻¹ when the parachute opens. Consider next the value of v when t is arbitrarily large. For such a value of t, $8e^{-1.25t}$ would be arbitrarily small, so v would be very close to the value 12. The modelling interpretation of this is that eventually the speed becomes very close to a constant value, 12 m s^{-1} which will be maintained until the parachutist lands.

The steady speed which is approached by the parachutist (or anything else falling against air resistance) is called the **terminal velocity**. The parachute, of course, is designed to ensure that the terminal velocity is sufficiently low (12 m s^{-1} in the specific case we have looked at here) to give a reasonably gentle landing and avoid injury.

Now consider what happens as t increases from near zero. When t is near zero, the speed will be near 4 m s⁻¹. The amount being subtracted from 12, through the term $8e^{-1.25t}$, is close to 8 because $e^0 = 1$. As t increases the value of $8e^{-1.25t}$ decreases fairly rapidly at first and then more gradually until v is very nearly 12. This is sketched in Figure 10. In fact v is never equal to 12 but gets imperceptibly close as anyone would like as t increases. The value shown as a horizontal broken line in Figure 10 is called an **asymptotic limit** for v.



Figure 10: Graph of a parachutist's speed against time

The model concerned the approach of a parachutist's velocity to terminal velocity but the kind of behaviour portrayed by the resulting function is useful generally in modelling any **growth to a limit**. A general form of this type of growth-to-a-limit function is

$$y = a - be^{-kx} \qquad (C \le x \le D)$$

where a, b and k are positive constants (parameters) and C and D represent values of the independent variable between which the function is valid. We will now check on the properties of this general function. When $x = 0, y = a - be^0 = a - b$. As x increases the exponential factor e^{-kx} gets smaller, so y will increase from the value a - b but at an ever-decreasing rate. As be^{-kx} becomes very small,



y approaches the value a. This value represents the limit, towards which y grows. If a function of this general form was being used to create a model of population growth to a limit, then a would represent the limiting population, and a - b would represent the starting population.

There are three parameters, a, b, and k in the general form. Knowledge of the initial and limiting population only gives two pieces of information. A value for the population at some non-zero time is needed also to evaluate the third parameter k.

As an example we will obtain a function to describe a food-limited bacterial culture that has 300 cells when first counted, has 600 cells after 30 minutes but seems to have approached a limit of 4000 cells after 18 hours.

We start by assuming the general form of growth-to-a-limit function for the bacteria population, with time measured in hours

$$P = a - be^{-kt} \qquad (0 \le t \le 18).$$

When t = 0 (the start of counting), P = 300. Since the general form gives P = a - b when t = 0, this means that

$$a - b = 300.$$

The limit of P as t gets large, according to the general form $P = a - b^{-kt}$, is a, so a = 4000. From this and the value of a - b, we deduce that b = 3700. Finally, we use the information that P = 600 when t (measuring time in hours) = 0.5. Substitution in the general form gives

$$600 = 4000 - 3700e^{-0.5k}$$
$$3400 = 3700e^{-0.5k}$$
$$\frac{3400}{3700} = e^{-0.5k}$$

Taking natural logs of both sides:

$$\ln\left(\frac{3400}{3700}\right) = -0.5k$$
 so $k = -2\ln(\frac{34}{37}) = 0.1691$

Note, as a check, that k turns out to be positive as required for a growth-to-a-limit behaviour. Finally the required function may be written

$$P = 4000 - 3700e^{-0.1691t} \qquad (0 \le t \le 18).$$

As a check we should substitute t = 18 in this equation. The result is P = 3824 which is close to the required value of 4000.



Your solution

Find a function that could be used to model the growth of a population that has a value of 3000 when counts start, reaches a value of 6000 after 1 year but approaches a limit of 12000 after a period of 10 years.

(a) First find the modelling equation:

Answer

Start with

 $P = a - be^{-kt} \qquad (0 \le t \le 10).$

where P is the number of members of the population at time t years. The given data requires that a is 12000 and that a - b = 3000, so b = 9000.

The corresponding curve must pass through (t = 1, P = 6000) so

 $6000 = 12000 - 9000e^{-k}$

$$e^{-k} = \frac{12000 - 6000}{9000} = \frac{2}{3}$$
 so $e^{-kt} = (e^{-k})^t = \left(\frac{2}{3}\right)^t$ (using Rule 3b, Table 1, page 42)

So the population function is

$$P = 12000 - 9000 \left(\frac{2}{3}\right)^t \quad (0 \le t \le 10).$$

Note that P(10) according to this formula is approximately 11840, which is reasonably close to the required value of 12000.

(b) Now sketch this function:





5. Inverse square law decay



Inverse square law decay of electromagnetic power

Introduction

Engineers are concerned with using and intercepting many kinds of wave forms including electromagnetic, elastic and acoustic waves. In many situations the intensity of these signals decreases with the square of the distance. This is known as the **inverse square law**. The power received from a beacon antenna is expected to conform to the inverse square law with distance.

Problem in words

Check whether the data in the table below confirms that the measured power obeys this behaviour with distance.

Power received, W	0.393	0.092	0.042	0.021	0.013	0.008
Distance from antenna, m	1	2	3	4	5	6

Mathematical statement of problem

Represent power by P and distance by r. To show that the data fit the function $P = \frac{A}{r^2}$ where A is a constant, plot $\log(P)$ against $\log(r)$ (or plot the 'raw' data on log-log axes) and check

- (a) how close the resulting graph is to that of a straight line
- (b) how close the slope is to 2.

Mathematical analysis

The values corresponding to log(P) and log(r) are

$\log(P)$	-0.428	-1.041	-1.399	-1.653	-1.851	-2.012
$\log(r)$	0	0.301	0.499	0.602	0.694	0.778

These are plotted in Figure 11 and it is clear that they lie close to a straight line.



Figure 11

The slope of a line through the first and third points can be found from

$$\frac{-1.399 - (-0.428)}{0.499 - 0} = -2.035$$

The negative value means that the line slopes downwards for increasing r. It would have been possible to use any pair of points to obtain a suitable line but note that the last point is least 'in line' with the others. Taking logarithms of the equation $P = \frac{A}{r^n}$ gives $\log(P) = \log(A) - n\log(r)$

The inverse square law corresponds to n = 2. In this case the data yield $n = 2.035 \approx 2$. Where $\log(r) = 0$, $\log(P) = \log(A)$. This means that the intercept of the line with the $\log(P)$ axis gives the value of $\log(A) = -0.428$. So A = 10 - 0.428 = 0.393.

Interpretation

If the power decreases with distance according to the inverse square law, then the slope of the line should be -2. The calculated value of n = 2.035 is sufficiently close to confirm the inverse square law. The values of A and n calculated from the data imply that P varies with r according to

$$P = \frac{0.4}{r^2}$$

The slope of the line on a log-log plot is a little larger than -2. Moreover the points at 5 m and 6 m range fall below the line so there may be additional attenuation of the power with distance compared with predictions of the inverse square law.

Exercises

- 1. Sketch the graphs of (a) $y = e^t$ (b) $y = e^t + 3$ (c) $y = e^{-t}$ (d) $y = e^{-t} 1$
- 2. The figure below shows the graphs of $y = e^t$, $y = 2e^t$ and $y = e^{2t}$.



State in words how the graphs of $y = 2e^t$ and $y = e^{2t}$ relate to the graph of $y = e^t$. 3. The figures below show graphs of $y = -e^{-t}$, $y = 4 - e^{-t}$ and $y = 4 - 3e^{-t}$.



Use the above graphs to help you to sketch graphs of (a) $y = 5 - e^{-t}$ (b) $y = 5 - 2e^{-t}$

4. (a) The graph (a) in the figure below has an equation of the form

 $y = A + e^{-kt}$, where A and k are constants. What is the value of A?

- (b) The graph (b) below has an equation of the form $y = Ae^{kt}$ where A and k are constants. What is the value of A?
- (c) Write down a possible form of the equation of the exponential graph (c) giving numerical values to as many constants as possible.
- (d) Write down a possible form of the equation of the exponential graph (d) giving numerical values to as many constants as possible.





6. Logarithmic relationships

Experimental psychology is concerned with observing and measuring human response to various stimuli. In particular, sensations of light, colour, sound, taste, touch and muscular tension are produced when an external stimulus acts on the associated sense. A nineteenth century German, Ernst Weber, conducted experiments involving sensations of heat, light and sound and associated stimuli. Weber measured the response of subjects, in a laboratory setting, to input stimuli measured in terms of energy or some other physical attribute and discovered that:

- (1) No sensation is felt until the stimulus reaches a certain value, known as the threshold value.
- (2) After this threshold is reached an increase in stimulus produces an increase in sensation.
- (3) This increase in sensation occurs at a diminishing rate as the stimulus is increased.



- (a) Do Weber's results suggest a linear or non-linear relationship between sensation and stimulus? Sketch a graph of sensation against stimulus according to Weber's results.
- (b) Consider whether an exponential function or a growth-to-a-limit function might be an appropriate model.

Answer



(b) An exponential-type of growth is not appropriate for a model consistent with these experimental results, since we need a diminishing rate of growth in sensation as the stimulus increases. A growth-to-a-limit type of function is not appropriate since the data, at least over the range of Weber's experiments, do not suggest that there is a limit to the sensation with continuing increase in stimulus; only that the increase in sensation occurs more and more slowly.

A late nineteenth century German scientist, Gustav Fechner, studied Weber's results. Fechner suggested that an appropriate function modelling Weber's findings would be logarithmic. He suggested that the variation in sensation (S) with the stimulus input (P) is modelled by

 $S = A \log(P/T) \qquad (0 < T \le 1)$

where T represents the threshold of stimulus input below which there is no sensation and A is a constant. Note that when P = T, $\log(P/T) = \log(1) = 0$, so this function is consistent with item (1) of Weber's results. Recall also that \log means $\log arithm$ to base 10, so when P = 10T, $S = A \log(10) = A$. When P = 100T, $S = A \log(100) = 2A$. The logarithmic function predicts that a tenfold increase in the stimulus input from T to 10T will result in the same change in sensation as a further tenfold increase in stimulus input to 100T. Each tenfold change is stimulus results in a doubling of sensation. So, although sensation is predicted to increase with stimulus, the stimulus has to increase at a faster and faster rate (i.e. exponentially) to achieve a given change in sensation. These points are consistent with items (2) and (3) of Weber's findings. Fechner's suggestion, that the logarithmic function is an appropriate one for a model of the relationship between sensation and stimulus, seems reasonable. Note that the logarithmic function suggested by Weber is not defined for zero stimulus but we are only interested in the model at and above the threshold stimulus, i.e. for values of the logarithm equal to and above zero. Note also that the logarithmic function is useful for looking at changes in sensation relative to stimulus values other than the threshold stimulus. According to Rule 2a in Table 2 on page 42, Fechner's sensation function may be written

$$S = A \log(P/T) = A[\log(P) - \log(T)] \qquad (P \ge T > 0)$$

Suppose that the sensation has the value S_1 at P_1 and S_2 at P_2 , so that

$$S_1 = A[\log(P_1) - \log(T)] \qquad (P_1 \ge T > 0)$$

 and

$$S_2 = A[\log(P_2) - \log(T)]$$
 $(P_2 \ge T > 0).$

If we subtract the first of these two equations from the second, we get

$$S_2 - S_1 = A[\log(P_2) - \log(P_1)] = A\log(P_2/P_1),$$

where Rule 2a of Table 2 has been used again for the last step. According to this form of equation, the change in sensation between two stimuli values depends on the ratio of the stimuli values. We start with

$$S = A \log(P/T) \qquad (1 \ge T > 0).$$

Divide both sides by A:

$$\frac{S}{A} = \log \frac{P}{T} \qquad (1 \ge T > 0).$$

'Undo' the logarithm on both sides by raising 10 to the power of each side:

$$10^{S/A} = 10^{\log(P/T)} = \frac{P}{T}$$
 (1 \ge T > 0), using Rule 4b of Table 2

So $P = T \times 10^{S/A}$ $(1 \ge T > 0)$ which is an exponential relationship between stimulus and sensation.

A **logarithmic** relationship between sensation and stimulus therefore implies an **exponential** relationship between stimulus and sensation. The relationship may be written in two different forms with the variables playing opposite roles in the two functions.

The logarithmic relationship between sensation and stimulus is known as the Weber-Fechner Law of Sensation. The idea that a mathematical function could describe our sensations was startling when



first propounded. Indeed it may seem quite amazing to you now. Moreover it doesn't always work. Nevertheless the idea has been quite fruitful. Out of it has come much quantitative experimental psychology of interest to sound engineers. For example, it relates to the sensation of the loudness of sound. Sound level is expressed on a logarithmic scale. At a frequency of 1 kHz an increase of 10 dB corresponds to a doubling of loudness.



Given a relationship between y and x of the form $y = 3\log(\frac{x}{4})$ $(x \ge 4)$, find the relationship between x and y.

Your solution

Answer

One way of answering is to compare with the example preceding this task. We have y in place of S, x in place of P, 3 in place of A, 4 in place of T. So it is possible to write down immediately

 $x = 4 \times 10^{y/3} \qquad (y \ge 0)$

Alternatively we can manipulate the given expression algebraically.

Starting with $y = 3 \log(x/4)$, divide both sides by 3 to give $y/3 = \log(x/4)$.

Raise 10 to the power of each side to eliminate the log, so that $10^{y/3} = x/4$.

Multiply both sides by 4 and rearrange, to obtain $x = 4 \times 10^{y/3}$, as before.

The associated range is the result of the fact that $x \ge 4$, so $10^{y/3} \ge 1$, so y/3 > 0 which means y > 0.

Log-linear Graphs





Introduction

In this Section we employ our knowledge of logarithms to simplify plotting the relation between one variable and another. In particular we consider those situations in which one of the variables requires scaling because the range of its data values is very large in comparison to the range of the other variable.

We will only employ logarithms to base 10. To aid the plotting process we explain how log-linear graph paper is used. Unlike ordinary graph paper, one of the axes is scaled using logarithmic values instead of the values themselves. By this process, values which range from (say) 1 to 1,000,000 are scaled down to range over the values 0 to 6. We do not discuss log-log graphs, in which both data sets require scaling, as the reader will easily be able to adapt the technique described here to those situations.

	• be familiar with the laws of logarithms
Prerequisites	• have knowledge of logarithms to base 10
Before starting this Section you should	 be able to solve equations involving logarithms
Learning Outcomes	• decide when to use log-linear graph paper
On completion you should be able to	• use log-linear graph paper to analyse functions of the form $y = ka^{px}$



1. Logarithms and scaling

In this Section we shall work entirely with logarithms to base 10.

We are already familiar with a particular property of logarithms: $\log A^k = k \log A$.

Now, choosing A = 10 we see that: $\log 10^k = k \log 10 = k$.

The effect of taking a logarithm is to replace a power: 10^k (which could be very large) by the value of the exponent k. Thus a range of numbers extending from 1 to 1,000,000 say, can be transformed, by taking logarithms to base 10, into a range of numbers from 0 to 6. This approach is especially useful in the exercise of plotting one variable against another in which one of the variables has a wide range of values.



Estimate the value of y when x = 1.35.

Solution

If we attempt to plot these values on ordinary graph paper in which both vertical and horizontal scales are linear we find the large range in the y-values presents a problem. The values near the lower end are bunched together and interpolating to find the value of y when x = 1.35 is difficult.





To alleviate the scaling p	roblem	in E	Example	e 10 er	nploy	logarith	ms to	scale o	down
the avalues giving:	x	1	1.1	1.2	1.3	1.4	1.5	1.6	
the <i>y</i> -values, giving.	$\log y$	0	0.33	0.63	0.97	1.17	1.41	1.63	_

Plot these values and estimate the value of y when x = 1.35.





2. Log-linear graph paper

Ordinary graph paper has **linear** scales in both the horizontal (x) and vertical (y) directions. As we have seen, this can pose problems if the range of one of the variables, y say, is very large. One way round this is to take the **logarithm** of the y-values and re-plot on ordinary graph paper. Another common approach is to use **log-linear graph paper** in which the vertical scale is a non-linear **logarithmic scale**. Use of this special graph paper means that the original data can be plotted directly without the need to convert to logarithms which saves time and effort.

In log-linear graph paper the vertical axis is divided into a number of **cycles**. Each cycle corresponds to a jump in the data values by a factor of 10. For example, if the range of *y*-values extends from (say) 1 to 100 (or equivalently 10^0 to 10^2) then 2-cycle log-linear paper would be required. If the *y*-values extends from (say) 100 to 100,000 (or equivalently from 10^2 to 10^5) then 3-cycle log-linear paper would be used. Some other examples are given in Table 3:

y - values	$\log y$ values	no. of cycles
$1 \rightarrow 10$	$0 \rightarrow 1$	1
$1 \rightarrow 100$	$0 \rightarrow 2$	2
$10 \rightarrow 10,000$	$1 \rightarrow 4$	3
$\frac{1}{10} \rightarrow 100$	$-1 \rightarrow 2$	3

Table 3

An example of 2-cycle log-linear graph paper is shown in Figure 14. We see that the horizontal scale is linear. The vertical scale is divided by lines denoted by $1,2,3,\ldots,10,20,30,\ldots,100$. In the first cycle each of the horizontal blocks (separated by a slightly thicker line) is also divided according to a log-linear scale; so, for example, in the range $1 \rightarrow 2$ we have 9 horizontal lines representing the values 1.1, 1.2, ..., 1.9. These subdivisions have been repeated (appropriately scaled) in blocks 2-3, 3-4, 4-5, 5-6, 6-7. The subdivisions have been omitted from blocks 7-8, 8-9, 9-10 for reasons of clarity. On this graph paper, we have noted the positions of A : (1,2), B : (1,23), C : (4,23), D : (6,2.5), E : (3,61).



Figure 14





On the 2-cycle log-linear graph paper (below) locate the positions of the points F:(2,21), G:(2,51), H:(5,3.5). [The correct positions are shown on the graph on next page.]









Example 12

It is thought that the relationship between two variables x, y is exponential

 $y = ka^x$

An experiment is performed and the following pairs of data values (x, y) were obtained

	x	1	2	3	4	5
-	y	5.9	12	26	49	96

Verify that the relation $y = ka^x$ is valid by plotting values on log-linear paper to obtain a set of points lying on a straight line. Estimate the values of k, a.

Solution

First we rearrange the relation $y = ka^x$ by taking logarithms (to base 10).

 $\log y = \log(ka^x) = \log k + x \log a$ · · .

So, if we define a new variable $Y \equiv \log y$ then the relationship between Y and x will be linear its graph (on log-linear paper) should be a straight line. The vertical intercept of this line is $\log k$ and the gradient of the line is $\log a$. Each of these can be obtained from the graph and the values of a, k inferred.

When using log-linear graphs, the reader should keep in mind that, on the vertical axis, the values are not as written but the logarithms of those values.

We have plotted the points and drawn a straight line (as best we can) through them - see Figure 15. (We will see in a later Workbook (HELM 31) how we might improve on this subjective approach to fitting straight lines to data points). The line intersects the vertical axis at a value log(3.13) and the gradient of the line is

$$\frac{\log 96 - \log 3.13}{5 - 0} = \frac{\log(96/3.13)}{5} = \frac{\log 30.67}{5} = 0.297$$

But the intercept is $\log k$ so

implying k = 3.13 $\log k = \log 3.13$

and the gradient is $\log a$ so

implying $a = 10^{0.297} = 1.98$ $\log a = 0.297$

We conclude that the relation between the x, y variables is well modelled by the

relation $y = 3.13(1.98)^x$. If the points did not lie more-or-less on a straight line then we would conclude that the relationship was *not* of the form $y = ka^x$.



Figure 15





Using a log-linear graph estimate the values of k, a if it is assumed that $y = ka^{-2x}$ and the data values connecting x, y are:

x	-0.3	-0.2	-0.1	0.0	0.1	0.2	0.3
y	190	155	123	100	80	63	52

First take logs of the relation $y = ka^{-2x}$ and introduce an appropriate new variable:

Your solution $y = ka^{-2x}$ implies $\log y = \log(ka^{-2x}) =$ introduce Y =

 $\log y = \log k - 2x \log a$. Let $Y = \log y$ then $Y = \log k + x(-2 \log a)$. We therefore expect a linear relation between Y and x (i.e. on log-linear paper).

Now determine how many cycles are required in your log-linear paper:

Your solution

The range of values of y is 140; from 5.2×10 to 1.9×10^2 . So 2-cycle log-linear paper is needed.

Now plot the data values directly onto log-linear paper (supplied on the next page) and decide whether the relation $y = ka^{-2x}$ is acceptable:

Your solution

It is acceptable. On plotting the points a straight line fits the data well which is what we expect from $Y = \log k + x(-2\log a)$.

Now, using knowledge of the intercept and the gradient, find the values of k, a:

Your solution

See the graph two pages further on. $k \approx 94$ (intercept on x = 0 line). The gradient is $\frac{\log 235 - \log 52}{-0.4 - 0.3} = -\frac{\log(235/52)}{0.7} = -\frac{0.655}{0.7} = -0.935$ But the gradient is $-2\log a$. Thus $-2\log a = -0.935$ which implies $a = 10^{0.468} = 2.93$



Your solution to Task on page 67





Answer to Task on page 67

Use the log-linear graph sheets supplied on the following pages for these Exercises.

Exercises

1. Estimate the values of k and a if $y = ka^x$ represents the following set of data values:

x	0.5	1	2	3	4
y	5.93	8.8	19.36	42.59	93.70

2. Estimate the values of k and a if the relation $y = k(a)^{-x}$ is a good representation for the data values:

x	2	2.5	3	3.5	4
y	7.9	3.6	1.6	0.7	0.3

Answers

1. $k \approx 4$ $a \approx 2.2$

2. $k \approx 200$ $a \approx 5$










NOTES

Index for Workbook 6

Asymptotic limit	50
Arrhenius' law	32-34
Base3,	23-26
Chemical reaction	32-34
Cosh	13
Coth	17
Cosech	17
Electromagnetic power	51
Equations	28
e	6-7
Exponent	3
Exponential function	_ 6-10
Exponential growth 8-9.	39. 48
Exponential decay10, 32,	45-48
Functions - even and odd	12
Hyperbolic functions	11-18
Index laws	22
Inverse square law	51
Kinetics	32
Kinetic energy	32
Linearisation	41-44
Log-linear graph paper	61
Log-linear plot 41-43,	58-73

50	Limit	_ 6, 48-50
32-34	Logarithm	19-26
0.00.00	Logarithm laws	22
3, 23-26	Logarithmic function	27-37
32-34	Natural logarithms	23
13		
17	Parameter	41
17	Population growth	50
51	Reverberation	46
28		
6-7	Scaling	59
3	Sinh	13
6-10	Sech	17
8-9, 39, 48	Solving equations	28-37
, 32, 45-48	Sound intensity	46
12	Tanh	17
11-18	Terminal velocity	48
22	EXERCISES	
51	10, 18, 26, 37, 53, 70	
30	ENGINEERING EXAMPLES	
32	1 Arrhenius' law	32
52	2 Exponential decay of sound	
41-44	intensity	46
61	3 Inverse square law decay of	
-43, 58-73	electromagnetic power	51

レ





HELM: Helping Engineers Learn Mathematics

http://helm.lboro.ac.uk